

# Bacteriophage T3 and Bacteriophage T7 Virus-Host Cell Interactions

DETLEV H. KRÜGER\* AND CORNELIA SCHROEDER

*Institute of Virology, Humboldt University, DDR-1040 Berlin, German Democratic Republic*

INTRODUCTION	9
GENERAL PROPERTIES	10
The Viruses and Their Growth Cycle	10
Restriction Analysis of Phage Deoxyribonucleic Acid (DNA)	10
EARLY VIRUS-CELL INTERACTIONS	11
Adsorption and DNA Injection	11
Nonclassical Modification of Viruses	12
REGULATION OF GENE EXPRESSION	12
Transcription	12
Early transcription	12
Late transcription	14
Messenger Ribonucleic Acid Processing and Translation	17
Shutoff of Host and Class I and Class II Phage Functions	18
INTERACTIONS WITH THE DNA MODIFICATION AND RESTRICTION SYSTEM OF THE HOST: FUNCTIONS OF GENE 0.3	20
Adenosylmethionine Hydrolase	20
Ability to Overcome Classical Restriction ( <i>ocr</i> )	20
Effects on DNA Modification	21
Modes of Action of <i>sam</i> <sup>+</sup> and <i>ocr</i> <sup>+</sup> Gene Functions	22
Other Functions Ascribed to Gene 0.3	22
DNA REPLICATION, MATURATION, AND RECOMBINATION	23
Enzymology of DNA Synthesis	23
Breakdown of host DNA	23
Primary initiation	24
DNA polymerase	24
Ribonucleic acid primer synthesis	24
DNA-unwinding protein and DNA ligase	25
DNA-relaxing enzymes	25
Reconstituted in vitro systems	25
Structure of Replicating and Maturing DNA	26
DNA Recombination	27
PHAGE MORPHOGENESIS AND CELL LYSIS	28
T3 and T7 Assembly	28
In vitro packaging	28
In vivo morphogenesis	28
Gene 3.5 Product and True Lysis Enzyme	29
INFLUENCE OF HOST-SPECIFIC FACTORS ON VIRUS MULTIPLICATION	29
<i>tsn</i> Mutations of <i>Escherichia coli</i>	29
Interference by F Plasmids	30
P1 Prophage	30
Antibiotic Resistance Plasmids and Colicinogenic Plasmids	30
<i>Shigella sonnei</i> D <sub>2</sub> 371-48	31
SPECIAL VIRUS-VIRUS AND VIRUS-HOST INTERACTIONS	31
T3 as a Facultative Temperate Phage	31
<i>E. coli</i> Transfection	32
Problem of Heterologous and Homologous Superinfection Exclusion	32
Interactions with Eucaryotic Systems	33
LITERATURE CITED	34

## INTRODUCTION

The recognition in the 1940s that bacteriophages could help to elucidate the molecular basis of heredity (49) has initiated exceedingly successful development. Phages, and viruses in

general, are relatively simple systems and much easier to analyze than are complex living systems. The study of bacteriophages has contributed much to the science of virology, since most phenomena originally observed in phage-infected bacteria were later on also discovered in

the interactions between eucaryote viruses and their host cells. The substantial technical advance during the past years has made it possible to explore molecular events in complex eucaryote cells and has established the intrinsic equivalence of pro- and eucaryote molecular genetics as well as a number of differences between them. The accessibility of eucaryote systems to molecular biological research does not, however, warrant the conclusion that the days of bacteriophage are over. Bacteriophages remain important models for virus-cell interaction: current research on bacteriophages is devoted to such problems as the mechanisms of virus-effected killing of cells, the integration and excision of viral genes in and from host genomes, deoxyribonucleic acid (DNA) modification and restriction, the reconstitution of the whole process of DNA replication in vitro, and solving the puzzle of temporal gene expression; these topics are of supreme interest for animal virology, too. In addition, bacteriophages have gained great scientific and economic prominence as vehicles for experimental gene transfer.

Bacteriophages T3 and T7 belong to the classical T series of bacteriophages (94). Not only the traditional *Escherichia coli* laboratory strains but also certain strains of *Shigella sonnei* (140, 164), *Salmonella typhimurium* (43), *Klebsiella pneumoniae* (379, 424), and *Pasteurella* (165) are permissive hosts. However, all essential knowledge about T3 and T7 virus-host interactions and the molecular biology of T3 and T7 was acquired in the *E. coli* system.

In writing this review it was a great advantage that we could rely on the excellent reviews of Hausmann (164, 165), McCorquodale (306), and Studier (484, 489), and, while concentrating on new developments, we shall refer to the 1976 review of Hausmann (165) for most of the work published up to that time. Faced with an avalanche of new papers, we are aware of the fact that the wealth of information can no longer be adequately presented in a condensed review; there is by now enough material to fill a book exclusively dedicated to these phages, a book like the modern classic *The Bacteriophage Lambda* (181). Therefore, we were forced to select topics for thorough treatment at the expense of others which were only briefly summarized.

## GENERAL PROPERTIES

### The Viruses and Their Growth Cycle

The principal features of the T3 and T7 phages' structure and growth cycle have been thoroughly reviewed (164, 165, 306, 484, 489) and are briefly summarized here. The virion consists

of a polyhedral nucleocapsid 50 nm in diameter to which a simple, noncontractile tail 20 nm long is attached which is capable of adsorbing to bacterial cells. The genome ejected through this tail is a single molecule of linear double-stranded DNA with a molecular weight of about  $25 \times 10^6$ , equivalent to approximately 40,000 base pairs. (The molecular weight of T7 DNA has recently been determined with greater accuracy to be  $26.4 \times 10^6$  to  $26.5 \times 10^6$  [76, 482]. According to a recent estimation, T3 DNA contains 38.74 kilobases [17], which is equivalent to a molecular weight of  $25.6 \times 10^6$ .) The DNA consists of the usual four nucleotides, the guanine plus cytosine content being similar to that of the *E. coli* host cell. The DNA is terminally redundant, but not circularly permuted, and codes for about 30 proteins. Gene expression of the phages is relatively simple, thanks to the exclusive transcription of the parental genome(s) and only of the heavy DNA strand thereof, as well as to the strictly sequential reading of the genes in the order in which they are arranged on the phage chromosome.

Phage development involves a number of regulated steps effecting shutoff of host and early phage functions, the degradation of the host genome into nucleotides, and their highly economical reutilization in the rapid synthesis of phage progeny DNA.

On infection of typical permissive *E. coli* strains under standard conditions, lysis begins at 12 to 15 min (37°C) or 20 to 25 min (30°C) and an average of 200 progeny phage are released per cell.

T3 and T7 are closely related, and their DNA molecules hybridize extensively; however, there are several functionally important nonhomologous regions. One of the salient differences between the two phages is the coding of an adenosylmethionine hydrolase (SAMase) by gene 0.3 of T3, but not by T7 (see Adenosylmethionine Hydrolase). Another nonhomologous region lies within gene 1, affecting the molecular weight and the template specificity of the ribonucleic acid (RNA) polymerase (RNA nucleotidyltransferase) (see Transcription). Differences between the two phages also show up in gene 17, where they determine the nonidentical antigenicities and adsorption specificities of the tail fiber proteins (see Adsorption and DNA Injection).

### Restriction Analysis of Phage Deoxyribonucleic Acid (DNA)

The utility of sequence-specific restriction endonucleases for physical mapping and sequence analysis of genomes and for gene cloning (272, 327, 328, 394, 395, 466) has had its impact also

on the elucidation of the structure and function of the T3 and T7 phage genomes. After the initial *Hind*II digestion performed by Kelly and Smith (220), and extension of this work (201, 259), several other restriction endonucleases were introduced into the analysis of T7 DNA (50, 147, 148, 275, 276, 300, 302, 307, 355, 494). The most extensive compilation of restriction analyses of the T7 genome was published by Rosenberg et al. (405). T7 DNA bears no recognition sites for the enzymes *Bam*HI, *Eco*RI, *Hind*III, *Pst*I, *Pvu*I, *Sal*I, *Sma*I, *Sst*I, *Sst*II, and *Xho*I; one to six recognition sites exist for *Ava*I, *Ba*II, *Bc*II, *Bgl*II, *Bgl*III, *Bst*EII, *Bst*NI, *Cl*aI, *Dpn*II, *Kpn*I, *Pvu*II, *Xba*I, *Xho*II (405), and *Eco*RII (W. Wackernagel, personal communication); and the DNA is cut at several sites by *Acc*I, *Alu*I, *Ava*II, *Hae*II, *Hae*III, *Hga*I, *Hgi*AI, *Hha*I, *Hind*II, *Hin*fI, *Hpa*I, *Hpa*II, *Hph*I, *Mbo*II, *Mn*II, *Sau*96I, *Taq*I, *Tha*I (405), and *Bsu*R (148). After initial experiments with *Hap*II, *Hga*I, *Hin*HI (502), and *Hpa*I (490), a detailed restriction map of T3 DNA has now been established by Bailey et al. (17). T3 DNA possesses no recognition sites for *Bam*HI, *Bgl*II, *Eco*RI, *Pst*I, *Pvu*II, *Sal*I, *Sma*I, *Sst*II, and *Xho*I; one to six sites for *Ava*I, *Bgl*III, *Hind*III, *Kpn*I, and *Xba*I; and several sites for *Hpa*I, *Mbo*I (17), *Hap*II, and *Hin*HI (502). An alignment of the T3 and T7 restriction maps reveals substantial differences (17, 405). Since differences also exist between laboratory strains of the same phage species, Studier (490) recommends a comparison of restriction patterns of phage from various laboratories in order to test their identity.

Physical maps of the early T7 genome region were established (147, 276, 300, 494), and certain genome fragments were isolated to study their interaction with *E. coli* and T7 RNA polymerase and for hybridization mapping of phage messenger RNA (mRNA) (200, 212, 217, 300, 302). In addition, restriction fragments were used for sequence analysis of T7 promoters (304, 336, 338, 404, 456).

Cloning of DNA fragments with the aid of plasmid or virus vehicles has opened new perspectives for the investigation of the molecular biology of T3 and T7. Campbell et al. (50) cloned some late and early T7 genes on the plasmid pMB9; in addition, the origin of DNA replication (51, 356, 423a) and promoter regions were cloned (346, 347, 435). Recently, the T7 gene 0.3 (*ocr*<sup>+</sup>) has been inserted into a bacteriophage lambda vector (363). Tsujimoto and co-workers took advantage of the fact that T7 wild-type DNA lacks *Eco*RI recognition sites to study its recombination with DNA of an *Eco*RI-sensitive T7 mutant (341, 508-510).

T7 DNA has a lasting place in the history of

science, as it was the first target for a sequence-specific restriction endonuclease (*Hind*II) tested in 1970 by Smith and co-workers (220, 467). From then on, type II restriction enzymes were used as tools for molecular biology (272, 327, 394, 466). It is curious that the interaction of *Hind*II with T7 DNA is of no significance in vivo, since T7 does not even adsorb to the *Hind*II-producing bacterium *Haemophilus influenzae* R<sub>d</sub> (D. H. Krüger and M. Hartmann, unpublished data).

## EARLY VIRUS-CELL INTERACTIONS

### Adsorption and DNA Injection

The *Host range* of a virus is defined as comprising those hosts which permit a productive viral growth cycle. We shall use this term in the more restricted sense of cells to which the virus can adsorb. The ability of the virus to adsorb and the presence or absence of virus receptors on a cell surface constitute the first level of virus-host interaction on which the establishment of infection is decided. The physical chemistry of virus adsorption has been largely neglected by current research, so there is little progress to report. The receptors for T3 and T7 consist of lipopolysaccharide areas in the outer membrane of the cell wall (38, 271, 373, 521). The tail fiber protein encoded by gene 17 is the main determinant of phage adsorption specificity (206, 207; O. G. Issinger, Ph.D. thesis, University of Freiburg, Freiburg, Federal Republic of Germany, 1973).

T3 and T7 genes 17 display a high degree of homology (89), which explains the related host ranges of these phages. The described host range of T7 is, however, broader than that of T3, since T7-resistant strains of *E. coli* are also resistant to T3, but not necessarily vice versa (94, 148, 159). Recent publications of Studier (490) and Krüger et al. (246) describe the resistance of *E. coli* K-12 Wisconsin (W) strains to T3 as opposed to their permissiveness for T7 infection; T3 host range mutants (T3hw) were isolated which adsorb to *E. coli* W as well as T7 (246). Another distinction between T3 and T7 is that the latter adsorbs with a higher rate to *E. coli* B cells (171; H. Beier, Ph.D. thesis, University of Freiburg, Freiburg, Federal Republic of Germany, 1973).

Corresponding to the similarity of the phage receptors, antisera to T3 and T7 are cross-reactive (3, 172, 206, 246; Issinger, Ph.D. thesis). The adsorption properties and serologies of T3 and T7 seem to be multifactorially determined, since the T3hw mutants adsorb to *E. coli* W, like T7, but retain their typical T3 adsorption pattern on different *E. coli* B strains (246, 249). The T3hw

mutation also alters the serological specificity of the phage, rendering it intermediate between those of T3 and T7 (246).

Triggered by cell-virus receptor interaction, the phage particle injects its DNA into the cell, beginning with the genetic left end (357, 417), the transcription of which appears to be the precondition for the complete injection of the genome (544).

### Nonclassical Modification of Viruses

It has been known for a long time that the ability of a phage to grow on a certain host strain can be dependent on a modification of the phage received in the previous host cell (277). When the modification is disadvantageous, phage growth on the new host is restricted. The criterion distinguishing *in vivo* modification and restriction (M/R) from mutational effects is the demonstration of reversible, host-dependent changes in the efficiency of plating of the phage (37, 277). Up to now, host-controlled M/R of bacteriophages has always been connected to modification versus cleavage of phage DNA (see Interactions with the DNA Modification and Restriction System of the Host: Functions of Gene 0.3). Krüger and co-workers (247-250) proved that, in addition, a totally different mechanism of host-dependent M/R exists which influences the adsorption ability of the phages and is caused by protein modification. In contrast to the "classical" M/R of DNA, this phenomenon was termed "nonclassical" M/R (247-250).

Such a nonclassical M/R was observed when T7 and SAMase-negative T3 derivatives were passaged on different *E. coli* B strains (250). The efficiency of plating of the phage varies from 0.01 to 1.0, dependent on the host strain on which the phage were last grown. These genetically unaltered phage carrying different nonclassical modifications also differ antigenically (K. K. Gachechiladze, D. H. Krüger, N. S. Bal-

ardzhishvili, S. Hansen, H. A. Rosenthal, and T. G. Chanishvili, submitted for publication).

A second case of nonclassical M/R is the behavior of T3 towards *E. coli* W1655 (248, 249). Independent of the fact that T3hw host range mutants appear at a frequency of  $10^{-8}$  in the phage population (see Adsorption and DNA Injection), the efficiency of plating of T3 wild-type phage assumes values between  $10^{-7}$  and  $10^{-1}$ , depending on the preceding host. These remarkable differences in efficiency of plating can be attributed quantitatively to different adsorption values (248, 249).

To our knowledge, nonclassical modification of T3 and T7 bacterial viruses constitutes a novel phenomenon in virology. The improvement of adsorption to cells of the same type and the impairment of adsorption to other cell types caused by modification may be related to the long-disputed mechanisms of cell and tissue adaptation of animal viruses which cannot be reduced to mutant selection alone (see Fenner et al. [123] p. 317-318). In addition, nonclassical M/R could interfere with lysotyping of bacteria (247).

## REGULATION OF GENE EXPRESSION

### Transcription

Transcription of phages T3 and T7 proceeds in distinct phases (Fig. 1): early transcription (class I genes), performed by the host RNA polymerase, succeeded by late transcription (class II and class III genes), carried out by the phage-specified RNA polymerase (484).

**Early transcription.** Early transcription commences at the left end of the genome, where *E. coli* RNA polymerase recognizes three adjacent promoters, A1, A2, and A3, *in vivo* (112, 318). The reader is guided to the 1979 review of Rosenberg and Court (406) for thorough information about promoter recognition. All early T7




FIG. 1. Outline of T7 genome organization. Data were compiled from references 143, 144, 165, 217, 302, and 484. For phage-coded proteins, see Table 4.

promoters (Table 1), with the exception of one minor promoter (*D* or *A0*) set off transcription in left-to-right direction to copy the heavy strand of viral DNA (471, 499). Promoters for bacterial RNA polymerases have in common homologies in functionally important regions which are strongly conserved in evolution (136, 406, 523) (Table 2). The “-35 region” is implicated in initial recognition, and the “Pribnow box” (374, 375, 427) probably triggers local unwinding of the double strand (444). By elegant chemical DNA modification techniques, Siebenlist (457) and Siebenlist and Gilbert (458) identified the minimal unwound promoter *A3* sequence (Table 2) and the contact points of the -35 region and the Pribnow box with *E. coli* RNA polymerase. It has been pointed out that the *C* promoter has the lowest homology to an “average promoter” constructed from superposition of several known promoter sequences, despite the fact that it is relatively strong under certain in vitro conditions (see discussions by McConnell [304] and Pfeffer et al. [362]).

The properties of purified *E. coli* RNA polymerase as well as its interaction with DNA have been reviewed extensively (20, 68, 69, 136, 211, 262, 543). The concentrations and the specificity

of bacterial RNA polymerase depend on the physiological state of the cell (2, 73, 130, 219, 274, 329, 392, 506, 507, 543). Since the role of the minor T3 and T7 promoters has not been firmly established, it would be interesting to know whether variant *E. coli* RNA polymerases are capable of recognizing alternative promoters on phage DNA.

The binding of *E. coli* RNA polymerase to T7 DNA has been explored by electron microscopy (36, 74, 87, 88, 232, 370, 525) and with the aid of isolated restriction fragments (148, 200, 202, 276). Improvements of the technique enabled Koller et al. (231) to reliably localize the minor promoters *B* through *E* (Table 1). These authors summarized and unified previously published data on the location of early T7 promoters. Remaining differences, e.g., seven binding sites identified by Koller et al. (231) as opposed to eight found by Hinkle and Chamberlin (190), may be reconciled by the work of Kadesch et al. (212, 213) proving the existence of tight-binding, promoter-like polymerase-DNA complexes distributed nonrandomly over the DNA molecule. The relative “strengths” of class I promoters estimated by three independent methods rise in the sequence  $A0 \ll A3 \leq A2 < A1$  (536).

The following steps precede initiation of mRNA synthesis: binding of RNA polymerase holoenzyme at “nonspecific” sites which might serve as storage stretches (426), diffusion to and recognition of promoters, conformational change of the enzyme, and melting of about 10 base pairs (87, 383, 457) of the promoter. At this stage the complex has become relatively stable to rifampin, heparin, polyinosinate, and denatured DNA (20). The rate of attack by rifampin (471) or heparin (362) depends on polymerase conformation, which, in turn, is determined by the structure of the respective promoter. The sensitivity of such initiation complexes in vitro does not correspond to the strength of the promoter (362). The actual initiation reaction is the formation of a dinucleoside tetraphosphate. Dinucleoside monophosphates stimulate rapid RNA synthesis in vitro (87, 289, 318, 339), which,

TABLE 1. Positions of promoters for *E. coli* RNA polymerase on T7 DNA

Promoter <sup>a</sup>	Position (% of genome) <sup>b</sup>	Preceding gene	Copied DNA strand <sup>c</sup>
<i>D</i> (= <i>A0</i> )	0.581		L
<i>A1</i>	1.221	0.3	H
<i>A2</i>	1.539	0.3	H
<i>A3</i>	1.834	0.3	H
<i>B</i>	3.728	0.7	H
<i>C</i>	7.821	1	H
<i>E</i>	92.123	19	H

<sup>a</sup> Designation according to Stahl and Chamberlin (471) and Studier et al. (494). These promoters are numbered I through VII by Koller et al. (231).

<sup>b</sup> Data from Koller et al. (231) determined electron microscopically. Errors estimated by these authors are 0.056 to 0.22% of genome length.

<sup>c</sup> L, Light; H, heavy.

TABLE 2. Sequences of promoters for *E. coli* RNA polymerase on phage T7 DNA<sup>a</sup>

Pro-moter	Sequence (sense strand, 5' → 3')
<i>A1</i>	AAAAGAGTA[TTGAC]TTAAAGTC—TAAC CTA TAGGATACTTACAGCCATCGAGAGGGACACGGCG
<i>A2</i>	AAACAGGTA[TTGAC]AACATGAAGTAAC ATG CAGTAAGATA CAAATCGCTAGGTAACACTAGCAG
<i>A3</i>	ACAAAAC <sup>C</sup> [TTGAC]AACATGAAGTAA—ACA CGGTACGATGTACCAATGAAACGACAGTGAGTC
<i>C</i>	ATAAGCAAC <sup>C</sup> [TTGAC]GCAATGTTA ATGG GC—TGA TAGTCT—TATCTTA

<sup>a</sup> Promoter sequences were compiled from references 374 (*A3*), 456 (*A1*, *A2*, and *A3*), and 304 (*C*). A or G indicates the first copied nucleotide; the solid box is Pribnow box, and the dotted box encloses the -35 region. Indicated homologies correspond to average promoter sequences (304, 404, 406, 456). Dashes (—) in *A1*, *A3*, and *C* were inserted to achieve exact superposition of homologous regions. The wavy line indicates the minimal unwound sequence (457).

however, does not initiate at the *in vivo* point (see references in the footnote to Table 2). Using purified *E. coli* RNA polymerase plus adenosine 5'-triphosphate and uridine 5'-triphosphate, Nierman and Chamberlin (330) succeeded in synthesizing T7 mRNA correctly initiated at the A1 promoter with pppApU, the natural 5' terminus of A1 mRNA (241).

In the elongation complex the DNA region protected by *E. coli* RNA polymerase is shorter (about 26 nucleotides in the coding strand [401]) than in the promoter complex (about 40 nucleotides [374, 375]).

Five early transcripts are made *in vivo* (39, 40, 112, 113, 177, 462, 486, 494, 497): 0.3, 0.7, 1.0, 1.1, and 1.3 mRNA (Fig. 2). The synthesis of T3 and T7 early messengers terminates either in a rho-factor-independent manner (226) at the major termination site TE1, located at 18.9% (376, 494) on the T7 genome, or at terminator sites behind transcripts 0.3, 0.7, and 1.0 in a rho-dependent process, as shown *in vitro* (84, 86) as well as *in vivo* (177). Adhya et al. (6) have identified two rho-dependent termination sites at 8% (after gene 0.7) and 15% (after gene 1) of T7 DNA. For details on mRNA termination, we refer the reader to the review of Adhya and Gottesmann (5). Reinitiation (318) at the beginnings of genes 0.7 and 1.0 (86)—supported by the existence of promoters at these positions (Table 1 and Fig. 2)—could contribute to the pattern of the gene expression.

The major terminator TE1 brings most of the transcribing complexes of T7 to a halt *in vivo* and *in vitro* (226, 318, 484). The same applies for T3 early termination *in vivo* (111, 226), but

apparently not *in vitro* (88, 111, 226). TE1 recognition *in vivo* is supposed to be promoted by the phosphorylation of the host polymerase  $\beta$  and  $\beta'$  subunits by the T7 0.7 gene product (gp0.7) (352, 368, 378, 410). Phosphorylation also renders the enzyme unable to reinitiate transcription of host and viral DNAs (546) (see Shutoff of Host and Class I and Class II Phage Functions). Read-through early mRNA terminates at site TE2 (30.1%) (318, 360, 462). Sequence analyses of the TE1 area of T7 DNA were recently published (116, 376).

**Late transcription.** Late transcription begins about 4 min (37°C) or 6 min (30°C) post-infection. Besides genes 1.1 to 1.3, the overlap region between class I and class II genes, the latter include genes 1.7 to 10. The first class II gene, immediately following TE1, has only been identified by DNA sequence analysis and precedes gene 1.7 (34, 116). Genes 7 and 10 are transcribed from both class II and class III promoters (303) (Fig. 1). By 1979 the primary structures of 10 late T7 promoters were elucidated (Table 3). These structures are shorter and mutually more homologous than promoters for bacterial polymerases (see Table 2). Regions of hyphenated twofold symmetry discovered in the 70% promoter by Oakley and Coleman (336) occur in all sequenced promoters and may contribute to promoter strength, since they are less pronounced in the class II promoters (Table 3). Class II promoters appear weak *in vitro* (22, 143, 144, 332, 463) but not *in vivo* (302, 346). Effective and specific transcription from class II promoters *in vitro* can only be accomplished by using restriction fragments lacking class III pro-




FIG. 2. Early region of phage T7 and its expression. Data compiled from references 34, 108, 112, 116, 217, 231, 276, 336, 354, 356, 398, 407, 418, 462, 491, 493, 494, and 503.

TABLE 3. Positions and sequences of promoters for T7 RNA polymerase on T7 DNA

Class	Position (% of genome)	mRNA species in vitro <sup>a</sup>	Sequence (sense strand, 5' → 3') <sup>b</sup>	Reference(s)
II	14.55	VII	CAATACGACTCACTATAGAGGGA	338, 355, 404, 494
II	14.75	VII	TAATACGACTCACTATAGGAGAA	355, 418
II	15.95	VII (?)	TAATACGACTCAGTATAGGGACA	116, 355
II	19.36	VIII	TAATACGACTCACTAAAGGAGGT	34, 116
II	19.65	VIII	TAATACGACTCACTAAAGGAGAC	34, 116
II	33.3	II <sub>b</sub> (?) (in vivo)	AATACGACTCACTAAAGGAXXC	Carter et al. <sup>c</sup>
II	34.75	II <sub>b</sub> (?) (in vivo)	TATTCGACTCACTATAGGAGAT	Carter et al. <sup>c</sup>
III	46.5	III <sub>a</sub>	TAATACGACTCACTATAGGGAGA	404
III	55	IV	TAATACGACTCACTATAGGGAGA	404
III	57	V	TAATACGACTCACTATAGGGAGA	404
III	70	II (?)	ACATACGACTCACTATAGGGAGA	338
III	87	III <sub>b</sub>	TAATACGACTCACTATAGGGAGA	404

<sup>a</sup> mRNA's are designated according to McAllister and McCarron (302), Kassavetis and Chamberlin (217), and Pachl and Young (346).

<sup>b</sup> The wavy line indicates the unwound sequence (481). The position of the initiation nucleotide (A) is questionable (see references 32, 34, 355, 404, 481).

<sup>c</sup> A. D. Carter et al., Abstr. Miami Winter Symp., 12:103.

motors (217), indicating that a special mechanism exists in vivo to activate class II promoters.

Since the first class II promoters are located immediately adjacent to gene 1, coding for phage RNA polymerase (418) it is probable that the enzyme is still physically connected to the gene (via mRNA and the ribosome) when it initially binds to the promoter. This mechanism of self-recognition is a general regulatory principle operative in procaryotes to ensure specificity (444).

Rosa (404) calculates that no more than 24 nucleotides in advance of the initiation site are necessary for promoter function. In her opinion the adenine- and thymine-rich sequence between -13 and -20 might be analogous to the -35 region, and the sequence TATAGGGA might be analogous to the Pribnow box TATPuATPu of promoters for bacterial RNA polymerase.

T3 and T7 RNA polymerases are monomeric enzymes of 97,000 and 105,000 daltons, respectively (Table 4). Both enzymes are specific for homologous DNA, that of T3 more strictly so than the T7 enzyme (22, 24, 66, 109, 145, 170, 281-283, 422, 430; Beier, Ph.D. thesis). They recognize neither host polymerase-specific promoters nor terminators (143, 144, 346) and are not amenable to rho-factor action (66).

Like practically all nucleotidyltransferases, T7 RNA polymerase is a zinc metalloenzyme (77), and like most other early T7 enzymes, RNA polymerase is found associated with the cell membrane (120). The initiation of RNA synthesis by the phage enzyme (301) is basically similar to the mechanism described for *E. coli* RNA polymerase (330); however, the polymerase-DNA complexes are unstable until a short oligonucleotide has been synthesized (65, 301). Like

the bacterial RNA polymerase, the T7 enzyme binds to the coding strand and melts a 10-base-pair sequence of the promoter (479) (see Table 3). RNA synthesis catalyzed by the T3 and T7 enzymes is 5-10 times faster than that of the bacterial polymerase (283, 301).

The specificity of promoter recognition by T3 and T7 RNA polymerase has been studied from different angles. Stahl and Chamberlin (472) showed that modification of the minor DNA groove by base analogs prevents utilization of promoters by T7 RNA polymerase, but alterations in one strand affecting the major groove do not. The properties of hybrid RNA polymerases induced by recombinants of T3 and T7 DNA within gene 1 were analyzed by Hausmann, Beier, and co-workers (22, 24, 170). The lower molecular weight of the T3 enzyme results from one or two small deletions in the T3 gene 1 when aligned with T7 gene 1 (24). Template preference seems to be imparted by the region from 0.70 to 0.78 gene 1 length units. The relative activity on heterologous phage DNA was found to vary from 20 to 80% among different recombinant RNA polymerases, in comparison with 15% for T3 and 50% for T7 wild-type RNA polymerases, and could not be ascribed to a particular region of the enzyme (170). In any case gene 1 recombinants are only viable if the template specificity domain of the RNA polymerase matches a minimum of promoter sequences within the recombinant DNA (22).

Chain initiation by T3 or T7 RNA polymerase is highly specific: the 5' terminus of all in vitro-synthesized mRNA is pppGp(Gp)<sub>n</sub>A in the case of the T3 (283) or pppGpGp(Ap)<sub>n</sub> in the case of the T7 (32) enzyme. Termination of RNA synthesis, at least by T3 RNA polymerase, appears

TABLE 4. *T7- and T3-directed proteins*<sup>a</sup>

Protein (gp) no.	Name or function of protein	Mol wt of protein (reference)		Constitu- ent of the mature virion	Reference(s)
		T7	T3		
0.3	<i>ocr</i> protein ( <i>ocr</i> and SAMase for T3)	8,700	11,500	No	254, 487, 493
0.4	Unknown	6,600 <sup>b</sup>	ND <sup>c</sup>	No	110, 474, 494
0.5	Unknown	ND	ND	No	494
0.6	Unknown	ND	ND	No	494
0.7	Protein kinase	42,000	40,000	No	44, 369, 352, 378, 485, 486, 410, 493
1	RNA polymerase	105,000 <sup>d</sup>	97,000	No	24, 70, 109, 333, 493; Issinger <sup>e</sup>
1.1	Basic protein (DNA initiation?)	5,700 <sup>b</sup>	ND	No	418, 494
1.2	DNA initiation?	12,000 <sup>b</sup>	ND	No	418, 494
1.3	DNA ligase	40,000	37,000	No	293, 346, 484, 485, 493
1.7	Unknown	17,000	ND	No	484
2	Inhibitor protein	8,500	ND	No	96, 185, 267
3	Endonuclease I	13,500	ND	No	61, 484
3.5	DNA and RNA metabolism (formerly called "lysozyme")	17,000	ND	No	228, 303, 346, 460, 484
<i>dup?</i>	DNA-unwinding (binding) protein	31,000	ND	No	165, 345, 385, 438
4	Primase	66,000 58,000 <sup>f</sup>	ND	No	188, 233, 434, 192, 484
5	DNA polymerase subunit, 3'-5' exonuclease	87,000	ND	No	197, 346, 388, 484
6	Exonuclease	31,000	ND	No	346, 484, 539
7	Coat protein (infectivity)	14,700 (484) 50,000 (207)	55,000	Yes	207, 319, 400, 484
8	Head protein	67,000	85,000	Yes	207, 319, 346, 400, 484
9	Head assembly (scaffolding) protein	45,000 <sup>d</sup>	ND	No	207, 319, 346, 400, 484
10	Major head protein	38,000 <sup>f</sup>	50,000	Yes	207, 319, 346, 400, 484
	Minor head protein	45,000			
11	Tail protein	22,000 <sup>d</sup>	27,000	Yes	207, 319, 400, 484
12	Tail protein	92,000 <sup>d</sup>	100,000	Yes	207, 319, 400, 484
13	Coat protein (infectivity)	14,000	13,000	Yes	207, 319, 400, 452, 484
14	Head protein (core)	20,000 <sup>d</sup>	20,000	Yes	207, 319, 400, 452, 484
15	Head protein (core)	89,000 <sup>d</sup>	96,000	Yes	207, 319, 346, 400, 452, 484
16	Head protein (core)	170,000 <sup>d</sup>	180,000	Yes	207, 319, 346, 400, 452, 484
17	Tail protein (fibers), "serum blocking power protein"	65,000 (346) 76,000 (207, 484)	80,000	Yes	206, 207, 319, 346, 400, 484
17.5 ?	Lysis enzyme	ND	ND	?	320, 325
18	DNA maturation and packaging	ND	ND	No	319, 400, 484
19	DNA maturation and packaging	73,000	ND	No	319, 346, 400, 484
20	DNA packaging in lambda-lysogenic cells	ND	?	?	358

<sup>a</sup> Molecular weights are either the latest published data, averages of published data, or, when differences between published data were too great, both currently accepted values. Recently identified small phage proteins of unknown function (34, 494) are not included.

<sup>b</sup> Computed from published DNA sequence (238, 418).

<sup>c</sup> ND, Not determined.

<sup>d</sup> Average data.

<sup>e</sup> O. G. Issinger, Ph.D. thesis, University of Freiburg, Freiburg, Federal Republic of Germany.

<sup>f</sup> Possibly processed.



to occur with comparable precision independent of other protein factors, yielding the 3' sequence Gp(Up)<sub>5</sub>G<sub>OH</sub> (285).

T3 and T7 RNA polymerases are salt sensitive in vitro (20, 65, 66, 109, 281, 430). The salt sensitivity of late T3 transcription in vitro can be overcome by a specific interaction of *E. coli dnaB* protein with T3 DNA (S. Michel and B. Brux, Abstr. Commun. Annu. Meet. Biochem. Soc. GDR, 11th, Halle/Saale, German Democratic Republic, 1979, E14; S. Michel, personal communication). By the following criteria this interaction appears also essential to relieving the salt sensitivity of late transcription in vivo: *E. coli* CRT 266, a *dnaB* mutant producing a thermosensitive *dnaB* protein, is nonpermissive for T3 replication at high temperature (290). Under nonpermissive conditions, only the early T3 proteins are expressed, and temperature shift experiments indicate that the *dnaB* protein is required for the initiation of RNA chains by T3 RNA polymerase. RNA synthesis in toluenized T3-infected *dnaB* mutant cells is stimulated by salt at permissive temperatures, but inhibited by salt at restrictive temperatures. Added wild-type *dnaB* protein reverses salt inhibition of purified T3 RNA polymerase at 42°C, whereas the thermosensitive *dnaB* mutant protein fails to do so (Michel and Brux, Abstr. Commun. Annu. Meet. Biochem. Soc. GDR, 11th, Halle/Saale, 1979, E14). One could surmise an analogy between the interaction of T3 RNA polymerase with *E. coli dnaB* protein, on the one hand, and that of *E. coli* primase with *dnaB* protein, wherein they form a "mobile replication promoter" (10), on the other.

Class III transcription encompasses all genes to the right of gene 6 (see Fig. 1). In vitro T3 RNA polymerase synthesizes six (145), seven (22, 285), or eight major transcripts (64, 65, 284) which are translatable by in vitro translation systems to phage proteins (65). T7 RNA polymerase generates six (333) or seven (143, 302) major mRNA species in vitro which are all class III (22, 332, 345). Besides that, three minor transcripts, A, B, and C, are detected which extend from promoters between 45 and 60% to the 99% termination site (302). Similar large-size transcripts (I, II, and III) are also generated by T3 RNA polymerase in vitro (284).

Late transcription terminates at only two specific sites (*TL1* and *TL2*) in vivo and in vitro: approximately 55 and 100% in the case of T3 (22) and 61 plus 99% in the case of T7 (302) (see Fig. 1).

Pachl and Young (346) have recently determined the sizes and mRNA activities of T7 late in vivo transcripts (eight from class II and eight

from class III) in an in vitro translation system and compared their data with the previously reported (22, 143, 144, 332) activity of mRNA synthesized in vitro. T3 and T7 in vivo transcripts either are identical to the in vitro transcripts or seem to be derived from them by ribonuclease (RNase) III digestion (see Messenger Ribonucleic Acid Processing and Translation). Among the in vivo transcripts it is possible to distinguish several class II mRNA's (155, 345, 346); these mRNA's cannot be fractionated from T7 mRNA synthesized in vitro (143, 144, 332) (see above). In T7-infected RNase III-negative host cells, Hagen and Young (155, 156) have detected T7 messengers as large as  $4 \times 10^6$  to  $5 \times 10^6$  daltons coding for gene 3.5 (class II) protein, indicating that class II mRNA synthesis in vivo terminates behind gene 10, at the first (*TL1*) of the two unique termination sites (Fig. 1).

### Messenger Ribonucleic Acid Processing and Translation

The early region of the T3 or T7 genome is transcribed into a large polycistronic messenger (39, 40, 112, 113, 177, 462, 497) which in vivo is processed by the host RNase III to yield 2 mono- and 3 polycistronic mRNA molecules (Fig. 2). The work on RNase III action not only is fundamental for the comprehension of the molecular biology of T3 and T7 but also has pioneered the study of sequence-specific RNA cleavage in eucaryote systems (1, 397). Thus, several publications were dedicated to the mechanism of site-specific RNA processing of T3 and T7 RNAs by RNase III (107, 108, 114, 178, 240, 241, 398, 408, 409, 491). The recognition sequences of RNase III were identified which presumably adopt a hairpin conformation (110, 114, 241, 336, 398, 407-409, 491). In vitro it is possible to generate cuts at secondary sites by lowering the ionic strength, whereby, for instance, fragment F5 is cleaved off the 1.1 mRNA (Fig. 2). Secondary processing is, however, relatively ineffective in vivo (107, 398). It is surprising that, in a situation resembling a primitive version of that in eucaryotic cells, the 3' ends of processed mRNA are subsequently oligoadenylylated (241, 407). T3 RNA polymerase catalyzes polyadenylate synthesis in vitro (422), but its involvement in oligoadenylation of phage mRNA has not been proven.

Hercules et al. (178) postulated that processing of the polycistronic early T3 and T7 messengers was necessary for their effective translation. Other authors (108, 114, 115, 488, 533) do not generally confirm this opinion, but find that the efficiency of 0.3 protein synthesis is increased by

RNase III cleavage of polycistronic mRNA, and possibly RNase III processing represents an evolutionary advantage under suboptimal growth conditions (9, 115, 178). The normally about fourfold excess of 0.3 mRNA over the other four early mRNA's is due not only to RNA processing but also to the rho-factor-mediated termination mentioned above.

Steitz and Bryan (474) and Dunn et al. (110) have studied mRNA binding and initiation of protein synthesis on T7 0.3 mRNA. Phage T7 0.3 mRNA is approximately 600 nucleotides long and carries the information for two proteins: the 5' two-thirds of the messenger specify the gp0.3 (see Interactions with the DNA Modification and Restriction System of the Host: Functions of Gene 0.3), and the remaining part codes for a protein of unknown function referred to as gp0.4. gp0.3 is made in large amounts during T7 infection, gp0.4 is made in smaller amounts, and each of these proteins is synthesized from separate specific ribosome-binding and initiation sites on the 0.3 mRNA. The proximity of the ribosome-binding site for gp0.3 synthesis to the RNase III cleavage site on the left end of gene 0.3 (474) may explain why RNase III action removing a nontranslated 5' RNA sequence is required for efficient 0.3 translation (115, 491).

Recently Studier et al. (494) demonstrated that not only 0.3 mRNA but also the 0.7 and 1.1 mRNA's encode more than one protein (0.7 mRNA encoding gp0.5, gp0.6, and gp0.7 and 1.1 mRNA encoding gp1.1 and gp1.2). Thus, the early region of the T7 genome codes for nine proteins. The functions of gp0.3 (*ocr* protein), gp0.7 (protein kinase), gp1.0 (RNA polymerase), and gp1.3 (DNA ligase [polydeoxyribonucleotide synthetase]) are ascertained, whereas nothing is known about the functions of the 50- to 120-amino-acid proteins 0.4, 0.5, 0.6, 1.1, and 1.2. For eight of these proteins (but not yet for gp1.2) the existence of special ribosome-binding and initiation sites which are apparently used with different efficiencies has been established (110, 747, 494). There is no evidence of any proteins being specified by the region between the A promoters at the far left of the genome (see Table 1 and Fig. 2) and the beginning of gene 0.3 (494).

Not only the early transcripts but also polycistronic class II and class III late T3 and T7 mRNA's are subject to RNase III processing (114, 157, 284, 346).

#### Shutoff of Host and Class I and Class II Phage Functions

Gene expression of T3 or T7 phage is regulated mainly on the level of transcription (177, 196, 478, 489). The early (class I) gene products serve to create an environment favorable for

phage-specified syntheses. Then the class II proteins responsible for the synthesis and maturation of DNA appear. As soon as sufficient amounts of these proteins are present, class I and class II transcriptions are turned off to ensure that the bulk of amino acids provided by the host cell are converted into phage structural and maturation proteins (class III proteins) (see Fig. 1 and Table 4).

The times of appearance and the concentrations of phage proteins are in essence determined by the positions of the respective genes in relation to promoters and termination sites. This does not apply only to their expression in one of the three classes: the dosages of certain proteins required for a longer period (possibly in the 1.1 to 1.3 region?) or in a very high number (the major head protein coded by gene 10) are enhanced because their genes are localized in areas of overlapping transcription—they are transcribed from early and late (genes 1.1 and 1.3) or from class II and class III late promoters (genes 7 to 10). Concentration differences between mRNA's and proteins of one class are also maintained by overlapping transcription from alternative promoters of this class. Thus, due to intermittent termination, promoter-proximal genes are transcribed more often than promoter-distal genes. The economy of phage syntheses displays such a degree of perfection that apparently genes 18 and 19, coding for maturation proteins (see Phage Morphogenesis and Cell Lysis), are transcribed only for the short period between the initiation of class III transcription and the termination of class II transcription (472, 484).

The most important regulatory principle consists in the induction of a phage-coded RNA polymerase with a strict specificity for homologous DNA and concomitant inactivation of the host RNA polymerase. Two phage proteins are involved in the inactivation of *E. coli* polymerase, first of all the T7 seryl-threonyl protein kinase (adenosine-5'-triphosphate-protein phosphotransferase) specified by gene 0.7 (352, 369, 378), which phosphorylates the  $\beta$  and  $\beta'$  subunits of *E. coli* RNA polymerase (546). The reason why the phosphorylated host enzyme becomes less active is not quite clear and cannot be explained satisfactorily by the recognition of the terminator TE1 by the modified polymerase (369) and its inability to reinitiate RNA synthesis (546). Recent studies by Hesselbach and Nakada (184) and McAllister and Barrett (298) confirm the role of the T7 protein kinase in the shutoff of *E. coli* RNA polymerase-dependent transcription. It is probable that the early T3 function capable of altering the  $\beta$  subunit of *E. coli* RNA polymerase and reducing polymerase

activity to 25% (100) is analogous to the T7 gene 0.7 function (despite the fact that these authors could not detect a corresponding activity in T7-infected cells).

gp0.7 phosphorylates not only the host's RNA polymerase but also other proteins in the infected cell (377), including itself in a self-inactivating reaction (352). Rahmsdorf et al. state that their discovery of T7 kinase is the first demonstration of a seryl protein kinase—a well-known enzyme activity in eucaryotes—in a procaryotic cell (378).

A much stronger effect on *E. coli* RNA polymerase is, however, exerted by an inhibitor protein coded for by gene 2 (184). This inhibitor protein (see Table 4) binds physically to RNA polymerase holoenzyme, forming a 1:1 complex (96, 185, 267). The sensitivity of the wild-type RNA polymerase is apparently dependent on the  $\sigma$  subunit, but its sensitivity can be abolished by *tsnB* and BR3 mutations affecting the  $\beta$  subunit of the enzyme (96) (see *tsn* Mutations of *Escherichia coli*). It is not clear whether a connection exists to the loss of the  $\sigma$  factor by *E. coli* RNA polymerase holoenzyme after T7 infection reported by Khourgess and co-workers (33, 548). gp2 has no effect on phage RNA polymerase (183–186, 267, 368, 369). An analogous late protein which blocks the initiation of RNA synthesis by *E. coli* RNA polymerase was also described for T3 (279, 280) and is probably T3 gp2.

In contrast to the 0.7 protein kinase, gp2 is essential for phage development (484). The exceeding complexity of gp2 functions (this protein is also required during DNA maturation [see DNA Replication, Maturation, and Recombination]) have made it difficult to pinpoint the essential interaction(s). It is clear, though, that binding of gp2 to *E. coli* RNA polymerase is essential, since the above-mentioned *tsnB* mutants (67) do not support the growth of T7 phage (96).

The shutoff of host and early phage protein syntheses is not entirely explained by the mechanisms described above. Two further processes may be involved in the early-to-late switch: translational discrimination in favor of late mRNA and functional instability of early mRNA's. Some authors (441, 536–538) have estimated the functional half-life of early mRNA to be only a few minutes, and, therefore, when early transcription is shut down, the cessation of early protein synthesis is inevitable. Results implicating an additional phage-directed control mechanism (protein) preventing the translation of early phage mRNA (179, 446) have been contradicted (532). The situation is complicated by the fact that Young and co-workers report longer mRNA half-lives (196, 346, 478, 479) than

those reported by Yamada's group (536–538) and postulate a translational control at least for the expression of gene 0.3 (478). Results of Strome and Young (480) do indicate a translational discrimination against 0.3 mRNA towards the end of the infectious cycle, which, however, is not caused by a phage-specific control protein but simply by the excess of late mRNA. Hercules et al. (177) confirmed the rapid functional inactivation found by Yamada et al. (536–538) and could not trace any translational discrimination among the individual early mRNA species. However, significant differences in translational utilization were reported for early mRNA's by Steitz and Bryan (474), Dunn et al. (110), and Studier et al. (494) and for late mRNA's by Niles and Condit (332). The patterns of in vitro protein synthesis, using late T7 mRNA's, show a striking correspondence to in vivo patterns (196, 332, 345, 346); hence, additional modes of "late" translational control in the infected cell probably do not exist.

The predominance of class III protein synthesis over class II protein synthesis (484) might in part result from a faster functional inactivation of class II mRNA's, though data on this topic are scarce (154). McAllister and Wu (303) have now demonstrated that not only the synthesis of late proteins but also late transcription itself is temporally regulated: class II and class III mRNA's are synthesized at different times and in different amounts. This regulation is partially mediated by gp3.5, since in its absence the turn-off of class II transcription at 15 min postinfection (30°C) is not performed (303).

gp2 is also suspected to play a role in this switchover. It has been proposed that complexes of gp2 with *E. coli* RNA polymerase might bind to sites in the late genome region and inhibit progression of the phage-coded RNA polymerase (218, 471). This mechanism is reminiscent of the action of repressor-like polymerase-rifampin complexes on T7 mRNA synthesis in vitro (35, 218), and, in fact, one of the activities of gp2, namely, in DNA packaging in vitro (see below), can be replaced by rifampin (267, 344). In line with this evidence, the class II-type shutoff of gene 19 transcription (472, 484) might be accomplished by an inhibitory *E. coli* polymerase complex bound to promoter *E* (see Table 1 and Fig. 1).

In summary, the following mechanisms may, in principle, contribute to the observed concentration differences between individual phage proteins in the infected cell: (i) promoter utilization and termination (transcription rate), (ii) mRNA half-lives, (iii) translational utilization of individual messengers, and (iv) functional stability of the proteins themselves.

## INTERACTIONS WITH THE DNA MODIFICATION AND RESTRICTION SYSTEM OF THE HOST: FUNCTIONS OF GENE 0.3

### Adenosylmethionine Hydrolase

T3 encodes an enzyme, SAMase (EC 3.3.1.2), which hydrolyzes intracellular *S*-adenosylmethionine (SAM), yielding thiomethyladenosine and homoserine (131, 141, 142). The corresponding gene (*sam*<sup>+</sup>) is located at the left end of the genome (168) and was numbered gene 0.3 (493, 496). The molecular weight of the T3 protein is 11,500 (493), but a higher molecular weight, 17,000, has also been reported (469). T7 does not induce a SAMase, although a gene 0.3 is present (131, 141, 142, 254, 462, 470, 484, 487, 494).

SAM is the methyl donor for the transmethylation of macromolecules (DNA, RNA, protein); it is involved in the synthesis of polyamines and in other biosynthetic pathways (52, 511). SAM plays a key role in the processes of host-controlled DNA M/R; classical modification is based on DNA methylation, and, moreover, type I restriction enzymes require SAM as an allosteric effector for DNA binding or the stabilization of this interaction (11, 153, 504, 542).

Although Hausmann (161) isolated a number of T3*sam*<sup>-</sup> mutants, the biological functions of SAMase remained obscure for a long time, since *sam*<sup>+</sup> and *sam*<sup>-</sup> T3 derivatives behaved identically on a variety of host cells and under different growth conditions (164). In 1975 the biological consequences of SAMase action were recognized by Krüger et al. (243). In starved cells T3 wild-type phage becomes temperate, whereas T3*sam*<sup>-</sup> and T7 multiply normally. When, however, methylation of T3*sam*<sup>-</sup> phage DNA is prevented, the mutant phage also establishes a lysogenic infection (see T3 As a Facultative Temperate Phage). A second trait conferred by the possession of SAMase is the partial protection of T3 against in vivo restriction by *Eco*P1 endonuclease (243, 251), probably because the lack of SAM in the T3-infected cell prevents stable binding of the restriction endonuclease to phage DNA (see P1 Prophage).

The possibility of turning off all SAM-dependent processes with the aid of the *sam*<sup>+</sup> gene makes phage T3 a convenient tool to study the consequences of preventing DNA methylation, type I restriction, and other cellular SAM-dependent reactions (82, 141, 149, 227, 243, 288, 425). Doubtless, this is an interesting technique, but it must be taken into account that, for example, DNA M/R is counteracted not only by SAMase but also by a second function localized

in gene 0.3, the *ocr*<sup>+</sup> function (see Ability to Overcome Classical Restriction (*ocr*)).

For a long period T3 was believed to be the only phage to encode SAMase (164, 203). Meanwhile, it has been demonstrated that coliphage S<sub>D</sub>, *Serratia* phage IV, and *Klebsiella* phage 11 produce SAMase (239, 331).

### Ability to Overcome Classical Restriction (*ocr*)

T3 and T7 can be passaged through *E. coli* strains of different DNA host specificities (e.g., *E. coli* strains C, B, and K) without being phenotypically restricted (122). This property is not unique to T3 but is also displayed by T7 and T3*sam*<sup>-</sup> derivatives, which means that even in the absence of SAMase there is sufficient protection against DNA restriction, despite the fact that both T3 and T7 DNAs contain recognition sites for *Eco*B (121, 254).

Studier (487) and Studier and Movva (493) showed that T3 and T7 gene 0.3 mutants are phenotypically restricted by *E. coli* strains B and K and carry the respective modifications after passage over these strains. After this basic discovery, the protection mechanism was studied intensively. T3 specifies the SAMase (*sam*<sup>+</sup> gene function) as well as the ability to overcome classical restriction, which was termed *ocr*<sup>+</sup> by Krüger et al. (254). It was demonstrated (254) that *sam*<sup>+</sup> and *ocr*<sup>+</sup> are distinct functions of the T3 gene 0.3; T7 only possesses the *ocr*<sup>+</sup> function. The *ocr* mutants of T3 and T7 are subject to classical M/R via DNA methylation versus endonucleolytic cleavage (254). The effect of *ocr*<sup>+</sup> consists in actively turning off the cellular restriction enzyme to protect the unmodified recognition sites on the phage DNA against cleavage. This protection is, moreover, extended to any foreign DNA simultaneously introduced into the cell. When T3 or T7 phage are inactivated by ultraviolet radiation so that they can no longer destroy the cell but still are able to express gene 0.3, then it is possible to introduce plasmid DNA by conjugation (244) or transformation (387) without it being restricted. The recipient cells survive and are able to replicate themselves as well as the newly acquired plasmid (Fig. 3).

T3 and T7 are the first phages whose mechanisms of turning off cellular restriction enzymes were closely examined (see Modes of Action of *sam*<sup>+</sup> and *ocr*<sup>+</sup> Gene Functions). Protection is definitely exerted against type I restriction, and an effect against type II restriction (*Eco*RII) is indicated by preliminary results (Wackernagel, personal communication). Meanwhile, antire-




FIG. 3. Protection by T3 or T7 *ocr*<sup>+</sup> gene function of foreign plasmid DNA against restriction. Plasmid DNA, a common vector in gene transfer, is restricted in normal restriction-proficient recipient cells if it contains recognition sites for the appropriate restriction system (top). After preinfection of cells with ultraviolet (UV)-inactivated T3 or T7 phage still able to express the *ocr*<sup>+</sup> gene, the cellular restriction endonuclease (R) is blocked; subsequently introduced foreign plasmid DNA is not destroyed (bottom). (From references 244 and 387.)

striction functions of other phages have been described, e.g., for T5 (42, 90, 91) and *Bacillus subtilis* phage  $\phi$ NR2 (287). The situation is much more complex in the case of T-even phages, where in addition to the glucosylation of hydroxymethylcytosine, further protective functions exist (99, 270).

Recently, a function of restriction endonuclease *EcoK* in the in vivo repair of DNA cross-links in *E. coli* was reported (25) where the restriction enzyme acts not nucleolytically, but rather as a regulatory protein or by protecting single-stranded DNA against nuclease attack (26). The *ocr*<sup>+</sup> gene function does not diminish this effect of *EcoK* (A. A. Belogurov, G. B. Zavilgelsky, and D. H. Krüger, submitted for publication).

#### Effects on DNA Modification

The expression of SAMase during T3 infection causing a depletion of SAM explains why T3 DNA carries no methyl groups whatsoever (131, 141). In contrast, the DNA of the SAMase-negative phage T7 is methylated (131, 141). However, T7 DNA, too, is undermethylated compared with host cell DNA: the ratios of 6-meth-

ylaminopurine to adenine and of 5-methylcytosine to cytosine in *E. coli* DNA are about 30 to 50 times higher than those in T7 DNA (U. Günthert, Ph.D. thesis, University of Tübingen, Tübingen, Federal Republic of Germany, 1975). Two reasons for this have been recognized. First, the gene 0.3 *ocr*<sup>+</sup> function prevents not only host-controlled restriction but also modification (245). This is why T7 lacks host-specific (*hsp*) methylation, which is present in the DNA of *ocr*<sup>-</sup> mutants (245, 254, 487). The prevention of *hsp* methylation by the *ocr*<sup>+</sup> function is the explanation of the observation (121) that the DNA of T7 wild-type virus grown in *E. coli* B is susceptible in vitro to cleavage by purified restriction endonuclease *EcoB*.

Second, it was shown (102) that the *dam* methylase (deoxyadenosine methylase [291]) acts only weakly on T7 DNA in vivo: the recognition sequence 5'-GATC-3' for *MboI* (135), which happens to be identical with the sequence methylated by the *dam* enzyme (133, 160), is not protected against *MboI* cleavage in vitro. This result is supported by data proving that the *DpnI* endonuclease, which cuts at methylated 5'-GATC-3' sites, scarcely attacks T7 DNA in

vitro, whereas the complementary enzyme *DpnII*, which specifically cleaves the nonmethylated identical sequence, accepts T7 DNA as a substrate (257, 258). Considering that *E. coli dam* enzyme sensitizes DNA to the action of *DpnI* (515), one can deduce that during T7 DNA replication *dam* acts on very few, if any, of the *dam*-specific sites in T7 DNA. It is speculated that T7 may replicate faster than its DNA can be methylated by this enzyme (258), or, more probable, that the *dam* protein is an integral component of the *E. coli* DNA replication complex, but not of the T7 replication apparatus, and therefore has no access to phage DNA (102).

Unfortunately, the situation has been somewhat confused by a recently published paper of Auer et al. (15) in which it is stated that the gene 0.3 product (termed "M" protein by this group [see below]) has no influence on the methylation of T7 DNA. The mutants used in this study were merely selected for their inability to grow on a restriction-proficient *E. coli* host and are not phenotypically modified *in vivo* (15). Consequently, no difference in the DNA methylation of such mutants as compared with the wild type is to be expected. Of course, authentic gene 0.3 *ocr*<sup>-</sup> mutants undergo host-specific modification (245, 254, 487, 493). A comparison of the mutants used by this group (15, 193, 364, 366, 446) with the precisely mapped and well-characterized gene 0.3 mutants of Studier and co-workers (487, 493, 494) is called for.

#### Modes of Action of *sam*<sup>+</sup> and *ocr*<sup>+</sup> Gene Functions

In T3, *sam*<sup>+</sup> and *ocr*<sup>+</sup> represent distinctive functions of the 0.3 protein. T7 gp0.3 has only the *ocr*<sup>+</sup>, not the *sam*<sup>+</sup>, function. The two functions interfere with different steps in the interaction between DNA and type I restriction enzymes. By destruction of intracellular SAM, the *sam*<sup>+</sup> function ab initio suppresses DNA methylation. The activation of type I restriction enzymes and the stability of the enzyme-DNA complex, which both depend on SAM, are reduced in the SAM-depleted cell. The effect of the *ocr*<sup>+</sup> function is not connected with SAM hydrolysis and interferes with a later step of the interaction between restriction endonuclease and DNA, preventing cleavage as well as methylation (245, 251, 254; D. H. Krüger, Int. UNESCO/ICRO Symp., Szeged, Hungary, 1977). The independence of the *sam*<sup>+</sup> and *ocr*<sup>+</sup> activities has recently been confirmed by *in vitro* studies on purified gp0.3's from T3- and T7-infected cells (470). Thus, T3 gp0.3 appears to be at least bifunctional. The isolated 0.3 proteins of T3 and T7 bind directly to the restriction

endonucleases *EcoB* and *EcoK* and exert *ocr* activity in this way (470; F. W. Studier, personal communication).

Cellular restriction enzymes usually destroy foreign DNA immediately upon its entry into the cell (461). The *ocr*<sup>+</sup> function bestows complete resistance to endonucleolytic cleavage to the phage DNA (254), and this protection depends on active synthesis of gp0.3 in the infected cell (423; D. H. Krüger, unpublished data). We have already mentioned that gene 0.3 is located on the far left end of the genome, which is the first to be injected (357, 417) and expressed (168, 462, 487, 493, 497). Furthermore, T7 (and maybe also T3) 0.3 mRNA has an increased rate of synthesis and is more stable than the other early mRNA's (see Regulation of Gene Expression), which leads to an especially high rate of gp0.3 synthesis (177). But how can the *ocr*<sup>+</sup> function ensure instantaneous protection of the phage DNA when gp0.3 only becomes detectable at 2 min (37°C) in postinfection (168, 177)?

The delayed injection of T3 and T7 DNAs may be the "trick" by which the phages prevent exposure of DNA recognition sites before the appearance of gp0.3. Transcription of the leftmost DNA region must be the precondition for injection of the rest of the genome. This has been elegantly confirmed by experiments showing that upon blocking of *E. coli* RNA polymerase by rifampicin or streptolydigin, injection of T7 DNA is inhibited (544). The development of the antirestriction function of phage T5 follows a similar schedule (42, 90, 91).

The two gene functions *sam*<sup>+</sup> and *ocr*<sup>+</sup> make T3 and T7 interesting model viruses for resolving individual steps of DNA M/R and their modulation by virus functions. They could also be useful in elucidating the *in vivo* significance of sequence-specific DNA methylation and cutting processes for DNA replication, repair, recombination, and transcriptional control (12, 25, 48, 81, 92, 137, 194, 261, 382, 396, 512; T. A. Trautner, B. Pawlek, U. Günthert, U. Canosi, S. Jentsch, and M. Freund, Mol. Gen. Genet., in press; see also T3 as a Facultative Temperate Phage). It was discovered recently that SAM-dependent endonucleases corresponding to certain methylases also exist in eucaryotic cells (47).

#### Other Functions Ascribed to Gene 0.3

Ratner (380) found that T7, but not T3, gp0.3 binds to *E. coli* RNA polymerase. The nature of this interaction has not been studied. On the basis of this observation it was supposed for a certain time (183) that the inhibitor protein later identified as gene 2 product (184) was encoded by gene 0.3.

In 1974 Herrlich et al. (179) claimed to have localized the T7 gene coding for a translational repressor (see Shutoff of Host and Class I and Class II Phage Functions) on the left end of the T7 genome, between the early promoters and the 0.7 (kinase) gene. However, in the year thereafter Herrlich and co-workers (446) reported that the translational repressor was a late gene product and instead postulated that gene 0.3 encodes the so-called M (membrane) protein (446), which was supposed to make the *E. coli* membrane more permeable, allowing an efflux of potassium ions and creating an optimal ionic environment for the synthesis and function of phage-induced enzymes (366, 367). Herrlich and co-workers postulated that this drop in ionic strength was the basis for the following functions of the M protein (366, 446): (i) killing of the host cell, possibly via inhibition of host macromolecular syntheses; (ii) protection of infecting T7 DNA against host nucleases of the *recBC* and restricting types; (iii) mutual and superinfection exclusion of T3 and T7; and (iv) alteration of phospholipid synthesis after T7 infection, resulting in the appearance of new polar lipids. None of these assumptions has been substantiated (see Shutoff of Host and Class I and Class II Phage Functions, *E. coli* Transfection, and The Problem of Heterologous and Homologous Superinfection Immunity). The true function of gp0.3, the blocking of restriction enzymes, is exerted not via a reduction of ionic strength, but rather through direct binding of the gp0.3 to the restriction enzyme (470).

Even the significance of the potassium efflux from the cell due to M protein action has been questioned. First of all, efflux had only been observed at subnormal (1  $\mu$ M) potassium concentrations in the medium (366), and, furthermore, S. Michel and co-workers (personal communication) have observed no such passive K<sup>+</sup> loss from T3-infected cells. In addition, Michel and Brux (Abstr. Commun. Annu. Meet. Biochem. Soc. GDR, 11th, Halle/Saale, 1979, E14) (see Transcription) showed that, in the case of T3, transcription by phage RNA polymerase is carried out *in vivo* despite the presence of normal potassium concentrations, provided that a functional *E. coli dnaB* protein is present.

It is a fact that T7 gp0.3 is found associated with the cell membrane (120), but so are most other phage proteins, including even the major coat protein (gp10). McAllister and Wu (303) reported that neither deletions in the early T7 genome region nor a high concentration of monovalent cations (0.15 M KCl) had any effect on late T7 mRNA synthesis *in vivo*. Condit (79) observed a permeability change late in infection

(beginning 16 min postinfection at 30°C) which is probably caused by gp3.5 (see Gene 3.5 Product and True Lysis Enzyme).

## DNA REPLICATION, MATURATION, AND RECOMBINATION

Great progress has been made in the study of DNA replication, and several reviews have appeared to which we shall refer for general facts (8, 105, 132, 211, 236, 237, 314, 390, 459, 522, 545). DNA replication is initiated through a highly specific recognition step (for review, see references 8, 236, and 444), at a unique origin site(s), in some cases by incision of one of the parental DNA strands and in others by the synthesis of an RNA primer ("ori" RNA) from the origin sequence. DNA replication is catalyzed by a delicate multienzyme apparatus, the replisome (237, 522). Depending on the size and structure of the DNA template, different types of proteins are required for DNA strand separation (72, 211, 256). The replication fork opens up in advance of the growing "leading strand," which is probably synthesized continuously in 5' to 3' direction, whereas the "lagging strand" (3' to 5') must by necessity be synthesized in fragments, each initiating with a short RNA primer (343, 545).

DNA synthesis of double-stranded DNA phage proceeds either by a rolling circle mechanism or bidirectionally on linear or circular templates and produces characteristic intermediate DNA structures (105). The newly replicated DNA has to "mature," i.e., be processed, in order to yield native-sized progeny DNA (105, 390), and the ultimate stages of DNA maturation are coupled to phage morphogenesis.

Apart from the first round of replication, the synthesis of T7 DNA is spatially separated from the cellular region where parental T7 DNA is being transcribed. This might be achieved by different sites of membrane attachment of transcribing and replicating complexes (187). It is noteworthy that in the case of T7, DNA replication is not the precondition for late transcription (67, 484), as it is for T-even phage.

## Enzymology of DNA Synthesis

**Breakdown of host DNA.** About 5 min after T7 infection (37°C), *E. coli* DNA is released (probably aided by gp3.5 [460]) from a fast-sedimenting structure and subsequently disrupted into fragments ( $2 \times 10^6$  daltons) by gene 3-specified endonuclease. These fragments are totally digested to deoxynucleoside monophosphates by the gene 6 exonuclease (61, 222, 223, 359, 414). T7 and T3 induce no nucleotide-synthesizing enzymes of their own and are thus fully depend-

ent on the precursors acquired by host DNA degradation. The two phage-coded nucleases (gp3 and gp6) are also involved in DNA recombination and concatemer formation (see below), whereas other apparently T7-specified endonucleases (57, 348, 411) have not been assigned any definite role in T7 DNA metabolism.

**Primary initiation.** The origin of replication (*ori*) on T7 DNA was placed at 17% by electron microscopy (106) and has now been mapped precisely by Saito et al. (418) as a 129-base-pair sequence between 14.73 and 15.05 map units. It spans the right half of the 14.75 promoter, a 61-base-pair noncoding intercistronic region, and the first 14 codons of gene 1.1. The intercistronic sequence contains seven copies of the palindrome TTAA, which would allow the formation of hairpins, and there is a gene 4 protein (primase) site, GACCC, within. Saito et al. assume that at this position leading-strand synthesis is initiated by the primase (see below) and proceeds rightward until the first primase site on the opposite strand is exposed. This initial rightward movement might explain why eye-form intermediates at position 17 are observed. Marker rescue studies after ultraviolet irradiation had also suggested that the origin of replication is situated to the left of 17% (45, 46). This major origin initiates 70% of phage DNA synthesis (388). Deletion mutants lacking this origin are apparently viable (462), and minor origins have been identified (388, 418, 435). The predominant secondary origin is located at approximately 4% (418), near promoter *B*.

The proximity of *ori* to class II promoters has been noted (106, 229, 230, 390), and it was proposed that T7 RNA polymerase plays a role in primary initiation. This is supported by Hinkle's observation that thermoinactivation of a temperature-sensitive T7 RNA polymerase causes a cessation of phage DNA synthesis (189). The function of T7 RNA polymerase in initiation is probably the transcriptional activation of the *ori* region (see Fig. 2), a process described in the review of Kolter and Helinski (236). A facultative role of gp1.1 and gp1.2 in primary initiation is assumed (418).

**DNA polymerase.** T7 DNA polymerase (DNA nucleotidyltransferase, EC 2.7.7.7) has an intriguing structure. It consists of two subunits (230): the phage gp5 (150, 322, 340) plus the product of the *E. coli* *tsnC* gene (321), which was identified as thioredoxin (292). Thioredoxin was originally isolated as the cofactor of ribonucleoside-diphosphate reductase (265). It is a protein of 12,000 daltons (292) with a very high cysteine content. In relation to its function in T7 DNA replication, it may be important that thioredoxin can be phosphorylated at cysteine-32

(80, 365). Native T7 DNA polymerase is a dimer of these two subunits (7, 322). These may be separately purified from T7-infected cells (197) lacking either thioredoxin (*tsnC* [*trx*] mutants [195]) or gene 5 protein, and, on reconstitution, they yield functional T7 DNA polymerase (7, 198).

T7 DNA polymerase displays 3'-5' exonuclease activity (150, 340). gp5 itself is a single-strand-specific 3'-5' exonuclease possessing a deoxynucleoside triphosphate binding site (7, 198) but nevertheless unable to catalyze even a single polymerization step (198). The reconstituted enzyme, besides catalyzing DNA synthesis, carries a 3'-5' exonuclease activity for double-stranded DNA. It also has the single-stranded DNA 3'-5' exonuclease activity of gp5 (7, 198). The relationship between structure and function of T7 DNA polymerase is reminiscent of the fact that the phage T4 DNA polymerase *amb22* fragment (which lacks the cysteine-rich 20% COOH-terminal region) is devoid of polymerase activity but retains 3'-5' exonuclease activity (334, 335).

**Ribonucleic acid primer synthesis.** Nascent T7 DNA fragments complementary to the heavy strand are linked to RNA (313). Discontinuous synthesis of one of the strands has also been demonstrated by electron microscopy (528, 529) and hybridization studies (476). RNA primers are synthesized by a multifunctional phage enzyme, the gene 4-coded primase (432, 439), essential in vivo for lagging-strand synthesis (476, 529). In vivo the primers are two to six nucleotides long and composed mainly of adenosine and cytidine; the first deoxynucleotide is deoxyribosylthymine monophosphate (342, 447). Primer structure in vivo is more diverse than that in vitro, where predominantly pppACCA and pppACCC are made (188, 402, 403, 433, 434), which are also the most effective primers for subsequent DNA synthesis. T7 primase has been purified and characterized (192, 233, 433, 434, 439, 476). It occurs in a 66,000-molecular-weight species (434) and a 58,000-molecular-weight species (233) (see Table 4) which display identical enzyme activities in vitro.

T7 primase has several copurifying activities (188): (i) single-strand-DNA- and (in the presence of T7 DNA polymerase) also double-strand-DNA-dependent hydrolysis of (preferentially deoxy)nucleoside triphosphates (234), providing the energy for (ii) catalytic participation in DNA strand separation (188, 433) as a "helicase" (see references 72 and 256), (iii) catalysis of RNA primer synthesis (432-434, 439), and (iv) stimulation of primer extension by T7 DNA polymerase (188, 402).

The native enzyme is a dimer of two identical



subunits (188). The intimate functional cooperation of T7 primase and DNA polymerase is revealed by their mutual sensitivities to each other's inhibitors: (i) dideoxythymidine triphosphate interrupts DNA chain growth and concomitantly stops gp4-catalyzed (deoxy)nucleoside triphosphate hydrolysis, and (ii)  $\beta$ - $\gamma$ -methylenedeoxyribosylthymine triphosphate, a primase inhibitor, arrests DNA synthesis (234). Neither the three *E. coli* DNA polymerases nor phage T4 DNA polymerases are stimulated by T7 gp4 (434); thus, the interaction between T7 DNA polymerase and primase is highly specific. T7 primase can, however, cooperate with T3 DNA polymerase (188). In many respects the T7 primase-DNA polymerase enzyme pair is functionally equivalent to the *E. coli* *dnaB-dnaG* protein complex, acting as a "general priming system" for DNA replication (10, 237, 473).

RNA primers accumulate in T7-infected cells lacking gp6 and *E. coli* DNA polymerase I-associated 5'-3' exonuclease (454). Purified gp6 displays RNase H activity (455), so gp6 is presumably responsible for primer degradation in vivo.

**DNA-unwinding protein and DNA ligase.** Whereas T7 DNA polymerase and primase are essential phage functions, T7 also induces a non-essential DNA-unwinding protein (the corresponding *dup* gene has not been precisely mapped as yet). The protein was purified and shown to specifically stimulate T7 DNA polymerase (385, 386, 438). In contrast to the specificity of the T7 DNA-unwinding protein, the analogous *E. coli* protein stimulates *E. coli* DNA polymerases II and III (holoenzyme) and also T7 DNA polymerase, accounting for the fact that *dup* protein is dispensable during T7 reproduction (323, 324). The properties of DNA-unwinding proteins (also termed helix-destabilizing proteins) were recently reviewed (72).

T3 and T7 gene 1.3 encode an adenosine 5'-triphosphate-dependent DNA ligase (293, 493) which can be substituted in vivo by *E. coli* ligase, although the latter is nicotinamide adenine dinucleotide dependent (21, 293). One may ask why T7 (and T3) encode proteins which are functionally replaceable by host proteins. The explanation is probably that the DNA-unwinding protein is needed in stoichiometric amounts and that DNA ligase must seal DNA fragments at a multitude of sites simultaneously, also necessitating a high enzyme concentration.

**DNA-relaxing enzymes.** DNA gyrase (topoisomerase II), an enzyme with the capability of relaxing and introducing superhelical turns, is essential for DNA replication in *E. coli* (see reviews by Jovin [211] and Champoux [72]). In

addition to topoisomerase I ( $\omega$  protein), two types of DNA gyrase (topoisomerases II and II') are present in *E. coli* which differ functionally as well as in the size of the smaller subunit (41). This subunit is encoded by the *gyrB* (*cou* [coumermycin resistance]) gene of *E. coli* and the larger subunit is encoded by the *gyrA* (*nalA* [nalidixic acid resistance]) gene (72). Itoh and Tomizawa (208) showed that coumermycin inhibits T7 DNA replication in *E. coli* wild type and concomitantly less in coumermycin-resistant host cells, indicating that at least the *gyrB* (*cou*) gyrase subunit is involved in phage DNA replication. DeWynngaert and Hinkle (97) confirmed these in vivo results but showed that gyrase is not required in vitro. Coumermycin also inhibits late T7 transcription (97).

The requirement for topoisomerases during the replication of circular double-stranded DNA is evident; the apparent role of DNA gyrase during T7 phage DNA replication (208) was explained by the collapsed state of replicating T7 DNA in vivo (95) or by its membrane attachment (97, 208), either severely limiting free rotation of the linear double strands. The validity of this interpretation has recently been challenged by Kreuzer and Cozzarelli (242), according to whom nalidixic acid inhibits the growth of T7 phage, whereas heat inactivation of a thermosensitive *nalA* (*gyrA*) gyrase subunit does not. They infer that nalidixic acid blocks T7 growth by a corruption of the enzyme which, per se, is not necessary for T7 DNA synthesis. This corruption could, for instance, consist in trapping of a gyrase reaction intermediate (495) and would explain the dominance of drug susceptibility in vivo (242). In this connection it is remarkable that certain phage T3 DNA polymerase (gene 5) *ts* mutations confer a partial nalidixic acid resistance to T3 replication in vivo (443). In the light of Kreuzer and Cozzarelli's work, it is warranted to reevaluate the effects of coumermycin on T7 DNA replication.

**Reconstituted in vitro systems.** Attempts to reconstitute T7 DNA replication in vitro date back to 1973 (230), and partially reconstituted systems containing T7 DNA polymerase and primase with or without DNA-binding protein were described later on (191, 432). Replication forks formed in vitro closely resemble those formed in vivo (31, 432). Masker and Richardson (296, 297) developed an in vitro system including *E. coli* DNA polymerase I which produces mature-size T7 DNA capable of transfecting *E. coli* spheroplasts. Richardson et al. (388) have succeeded in reconstituting an enzyme system which performs primary initiation on intact duplex T7 DNA (see above) and carries out leading- and lagging-strand synthesis at the replica-

tion fork in a manner similar to that observed *in vivo*. Thus, the reactions occurring at the replication fork are now thoroughly understood (see Fig. 4). However, the role of the replicative intermediates described in Structure of Replicating and Maturing DNA is still not clear.

### Structure of Replicating and Maturing DNA

T3 and T7 DNAs possess terminal redundancies (391) which are essential for phage multiplication (22, 104). The length of the terminal

repetition of T7 DNA has now been estimated by independent methods to be 150 to 160 base pairs (103, 405). Due to unidirectional DNA polymerase movement, unit-size DNA has no means of replicating the 3'-terminal gaps which arise by excision of primer RNA. Watson (520) therefore postulated that linear duplex DNA must replicate via concatemers which form by hybridization of the unreplicated 3' redundant regions of newly synthesized DNA. To this day, Watson's postulate has not been conclusively proved or disproved.

The first round and at least the second round




FIG. 4. Stages of phage DNA replication. For explanations, see text. L, Light strand; H, heavy strand.

of T7 DNA replication begin at the *ori* site (see above) and proceed bidirectionally, creating first an eye and later on a Y-shaped intermediate structure (106, 528, 530); no circles are formed (530) (see Fig. 4).

Intracellular T7 DNA is associated with the membrane. (For technical details of studying phage DNA-membrane interactions, please consult the review of Siegel and Schaechter [459].) At a few minutes postinfection, parental T7 DNA is found associated with the inner cell membrane (also in the absence of phage protein synthesis) (58, 59, 347). The membrane complex contains phage DNA polymerase and actively synthesizes DNA (59); this DNA synthesis is just as ultraviolet sensitive as T7 *in vivo* DNA replication (526). Apparently, parental and progeny DNA molecules are bound to the membrane with different strengths (187). Parental DNA remains attached to the membrane throughout infection (187)—the attachment site is located within the early DNA region (175)—whereas replicating DNA is eventually released from the membrane (450). Parental DNA seems not to participate in concatemer formation (126). Since T7 DNA replication *in vitro* does not require membrane components, membrane association seems to serve an organizational role *in vivo*.

Concatemer DNA in T7-infected cells was initially observed by Hausmann (162), Carlson (53), Kelly and Thomas (221), Hausmann and LaRue (169), and Ihler and Thomas (204). Later during replication, there appear condensed coiled progeny DNA structures (30, 60, 187, 295, 350, 372, 440, 450, 476, 477, 510), which are also seen in T3-infected cells (129). This complex DNA was purified and characterized (260, 350). It contains about 100 phage genome equivalents and sediments at approximately 440S. In contrast to the compact folded *E. coli* genome (531), the T7 DNA complex is not held together by RNA, but contains hydrogen-bonded unit-size phage DNA (126, 440, 450, 477). Concatemer formation does not require recombination and may proceed when both the gene 3 and the gene 6 nucleases are simultaneously absent (126, 315). Under normal circumstances gp6 is essential for T7 (126, 169, 315) or T3 (539) concatemer formation. gp3 of T7 is not essential for *in vivo* and *in vitro* concatemer formation (295, 315, 415). The apparent requirement of gp3 for T3 DNA concatemerization probably reflects the polar effect of a gene 3 amber mutation on the expression of the *dup* gene. T3 DNA-unwinding protein may function in concatemer formation *in vitro* and also *in vivo* (129). T7 gp2 is required for the maintenance of concatemer structure (60).

The pathway of DNA replication and matu-

ration is conceived to be as follows: unit-size DNA → short concatemers → complex fast-sedimenting intermediate → short concatemers → unit-size DNA (350). Net T7 DNA synthesis is completed at 17 min postinfection at 30°C (260). The rapid, site-specific cleavage of concatemers into monomers is performed at the junctions between adjacent monomers, after the repair of gaps or protruding "whiskers" at these junctions (440). This is in accord with the above-mentioned postulate of Watson (520). The intimate connection between DNA maturation and packaging is discussed in Phage Morphogenesis and Cell Lysis.

### DNA Recombination

Molecular recombination is inherent to, but not necessarily essential for, T3 and T7 development and is coupled to the most active phase of DNA replication (294, 372). Recombination *in vivo* requires gp3, gp4, gp5, and gp6 (225, 371). Recombining DNA is fragmented before recombination (415), and gene 6 exonuclease is necessary for breakage and reunion of the fragments in parental-to-progeny DNA recombination (269, 315, 508). In the case of cloned DNA fragments, recombination into T7 DNA does not require gp4, gp5, and gp6 (51).

Biparental recombining T7 DNA molecules adopt characteristic intermediate structures *in vivo* (H- and Y-shaped) in a gp6-catalyzed reaction in the absence of gp3 (510). These branched structures are stabilized by gp5 (which plays a role other than gap filling) and are converted to linear recombinant molecules by the action of gene 3 endonuclease I (510). The same intermediates as *in vivo* arise in extracts of T7-infected cells (341).

Sadowski and Vetter (415) devised an *in vitro* assay which measures recombination of exogenous mature T7 DNA with endogenous T7 DNA present in extracts of infected cells. These extracts promoted recombination as well as packaging, so the recombinants could be assayed as plaque-forming units. This biological assay was complemented by a (less sensitive) physical assay (514) measuring the density shift of labeled exogenous phage DNA. It became clear that polar packaging (see Phage Morphogenesis and Cell Lysis) simulates asymmetric recombination in the biological assay (393), so Roeder and Sadowski (399) developed mutually independent and noninterfering *in vitro* T7 DNA recombination and packaging assays, proving that recombination is not inherent to *in vitro* packaging. This new recombination system, designed to promote recombination between exogenous DNA molecules, has permitted at least two pathways of *in vitro* recombination to be distin-

guished in appropriate mutant extracts: the endonuclease pathway, dependent on gp3, gp5, and either host or phage (gp1.3) ligase and inhibited by gp4 plus deoxynucleoside triphosphates, and the exonuclease pathway, requiring gp6, independent of ligase and gp3, gp4, and gp5, and stimulated by deoxynucleoside triphosphates. The wild-type *in vitro* pathway seems to combine features of both these separable pathways (399). As to enzyme requirements, the main difference between the *in vitro* and the *in vivo* reactions is the dispensability of gp4 and gp5 *in vitro*. Roeder and Sadowski (399) suggest that *in vivo* the physical separation of parental DNA molecules (bound to different membrane sites) may exclude biparental recombination so that only replicating DNA can recombine. On the other hand, according to the Radding-Meselson model (199), the products of genes 4 and 5 may be involved in DNA synthesis at nicks or gaps. Electron microscopy of recombining T7 DNA suggests that recombination is primed by an invasion of single-stranded DNA into single-stranded gaps at the replication fork (372).

*In vitro* recombination systems will probably soon allow the complete fractionation, reconstitution, and functional characterization of the individual recombination factors.

## PHAGE MORPHOGENESIS AND CELL LYSIS

### T3 and T7 Assembly

***In vitro* packaging.** *In vitro* packaging is an approach to resolve the steps of phage assembly and is also a useful procedure for assaying *in vitro*-manipulated DNA (see DNA Recombination). By heat-inactivating mature phage, Serwer (451) obtained a T7 DNA-nucleocapsid structure consisting of a unit length of T7 DNA to which an empty phage head is attached. Similar structures were seen in lysates of T7-infected cells. In the *in vivo* structure the attachment site was 7% from the genetic right end of the DNA; the detection of DNA-head complexes containing double-length DNA indicated that head binding is involved in the ultimate DNA maturation step (451).

Kerr and Sadowski (224) developed a combined *in vitro* DNA recombination and packaging assay. To study packaging uninfluenced by DNA recombination, Kuemmerle and Masker (255) introduced the use of gene 3, 5, and 6 triple mutants. Concatemeric DNA is packaged 10<sup>4</sup> times more efficiently than unit-size DNA (295), indicating that concatemer formation is a prerequisite for DNA packaging *in vivo*. *In vitro* packaging depends on the presence of the follow-

ing components (413): proheads (400); the proteins coded by genes 18 and 19, which are necessary for head filling (224, 400); and the tail proteins gp11, gp12, and gp17.

Fujisawa et al. (128, 319) developed an analogous *in vitro* packaging system for T3 DNA. Mature T3 DNA is converted to concatemers by the combined action of gp3 and gp6 and packaged into proheads in the presence of gp4, gp5, and gp19.

***In vivo* morphogenesis.** The basics of virus morphogenesis have been reviewed (55, 326). Since the 1976 review of Hausmann (165), a number of new papers dealing with T7 and T3 morphogenesis have appeared (117, 128, 207, 319, 400, 449, 452, 453, 534), of which we shall present a brief summary.

The proteins gp8, gp9, gp10, gp13, gp14, gp15, and gp16 assemble into proheads, and in the presence of gp18 and gp19 plus DNA, packaging is initiated. gp9 is a scaffolding protein necessary for the formation of the packaging intermediate but no longer present in the mature phage particle. Heads which have initiated, but not completed, DNA packaging are unstable and dissociate into DNA and empty heads. The final DNA maturation steps are synchronous with the completion of packaging. Somewhere between prohead formation and the completion of packaging, a host factor is active. Full heads do not contain gp18 and gp19.

gp11 and gp12 attach to full heads, forming the conical tail to which the tail fibers (gp17) are subsequently bound.

Mature phage heads consist mainly of gp10. In addition, gp14, gp15, gp16, and, probably, gp13 form a core which is surrounded by DNA and extends into the tail. This core structure might constitute an initiation complex during capsid assembly. gp8 is located at the junction of the phage head, core, and tail. gp7 and gp13 are not necessary for assembly, but they are necessary for the infectivity of the mature particle. All maturation steps proceed without demonstrable proteolytic cleavage.

For the proportions of the individual T7 structural proteins within the mature phage particle, see Hausmann (165).

The ultimate DNA maturation and packaging steps are blocked in T7 gene 2 mutants (96, 267, 344). gp2 is essential for concatemer processing (267, 344). This requirement appears to involve inhibition of *E. coli* RNA polymerase by gp2 (as described in Shutoff of Host and Class I and Class II Phage Functions), inasmuch as gp2 is replaceable by rifampin *in vivo* (344) and *in vitro* (267) and as *E. coli* *tsnB* mutants which induce a gp2-tolerant RNA polymerase also impose a

barrier on late T7 DNA replication (see *tsn* Mutations of *Escherichia coli*). On the basis of published data it is conceivable that an immobile gp2-RNA polymerase complex firmly attached to promoter *E* (92% on T7 DNA [see Table 1]) serves as the landing base for the phage prohead. This position corresponds exactly to that observed by Serwer for the binding of empty heads to T7 DNA (451). The contact point on the prohead might be gp10. If this were so, many isolated observations concerning the role of *E. coli* RNA polymerase and phage gp2 and gp10 would fall into place. Host mutants unable to support T7 DNA maturation, due to a disturbed interaction between host RNA polymerase and gp2 (96, 267), map in *rpoB*, the structural gene for the RNA polymerase  $\beta$  subunit. The *E. coli* Y49 mutant selected for its inability to plate T7 0.7 (protein kinase)-negative mutants, imposes a similar block on phage maturation (534, 535). It is probable that the Y49 mutation, like the BR3 mutation described by Studier, which is also nonpermissive for gene 0.7 mutants (485), maps in *rpoB* as well. An abortive packaging reaction is also observed in *S. sonnei sin* (see Influence of Host-Specific Factors on Virus Multiplication) which can be overcome by a mutation in T7 gene 10 (513). *sin* may be yet another *rpoB* mutation. If this explanation is correct, then there is in fact only one host factor involved in phage T7 maturation, the bacterial RNA polymerase.

### Gene 3.5 Product and True Lysis Enzyme

The role of the gp3.5 ("lysozyme"), an N-acetylmuramyl-L-alanine amidase, has been debated for several years (93, 127, 205, 210, 228, 484). This enzyme was initially thought to be responsible for cell lysis (180, 205). Studier (484) already pointed out that gp3.5 is expressed among the class II genes whose products function in DNA metabolism and, furthermore, that T7 gene 3.5 mutants are capable of inducing cell lysis. Jensen and Pryme (210) demonstrated that the standard test involving the liberation of radioactive material from diamino[ $^3\text{H}$ ]pimelic acid-labeled *E. coli* cell walls (445) is no proof of cytolytic potency in vivo and identified an additional amidase activity in T7-infected cells. The latter is responsible for cell lysis (210). Miyazaki et al. (320) and Blair et al. (cited in reference 25) showed that the true lysis enzyme is encoded by a class III gene in T3 and T7 located somewhere between genes 17 and 19 (gene 17.5?).

gp3.5 functions in DNA metabolism and probably promotes the release of host and newly made phage DNA and phage particles from the

cell membrane (460). It also seems to be engaged in the shutoff of class II mRNA synthesis (303).

Since the designation "lysozyme" is associated with active disruption of the cell wall and cell lysis, it would be sensible to abandon this name for the gene 3.5 product. To avoid confusion between gp3.5 and gp17.5, we propose to call the latter the lysis enzyme.

The lysis enzyme destroys the cell and liberates progeny phage particles. However, even in its absence, cell destruction is bound to occur eventually as the result of extensive phage-mediated cell damage (see Regulation of Gene Expression).

### INFLUENCE OF HOST-SPECIFIC FACTORS ON VIRUS MULTIPLICATION

The study of bacteria which have lost or attained the ability to support virus growth is often very useful for defining the role of certain bacterial elements in phage development and complements the investigation of phage mutants. In the course of this review we have mentioned a number of host proteins which are active during different stages of the phage growth cycle: the *E. coli* RNA polymerase and thioredoxin as the most important of these factors; the *dnaB* protein, which is essential for late T3 transcription; and an as yet ill-defined host factor possibly involved in a late morphogenesis step (534, 535). In the following section we will discuss the roles of certain host factors in greater detail.

#### *tsn* Mutations of *Escherichia coli*

The isolation of *E. coli tsnB* and *tsnC* mutants by Chamberlin (67) was of great importance for T7 research. The further study of the *tsnC* mutation led to the fundamental discovery that the *tsnC* gene is the structural gene for thioredoxin (*trx*) and that *E. coli* thioredoxin is a necessary constituent of T7 DNA polymerase (see Enzymology of DNA Synthesis).

*tsnB* mutations map within the structural gene for the  $\beta$  subunit of *E. coli* RNA polymerase (*rpoB*) (see review by Yura and Ishihama [543]). Investigations of *tsnB* mutants and of another *rpoB* mutant, *E. coli* BR3, which is unable to sustain the growth of T7 gene 0.7 mutants (485), proved that *E. coli* RNA polymerase is indispensable not only for early phage transcription but also for a late DNA maturation step (see Structure of Replicating and Maturing DNA) (96, 98, 267).

In an  $\text{F}^+$  background, nonpermissive for T7 reproduction (see below), the introduction of *strA* mutations (affecting ribosomal protein S12) allows T7 phage to overcome the  $\text{F}^+$  barrier. The

original, nonpermissive state can be fully restored by an additionally introduced *rpoB* (rifampin resistance [*rif*]) mutation (62, 63). A survey of *rpoB* mutants of different phenotypes as to their effects on phage replication would probably greatly further our understanding of T7 virus-host interaction.

### Interference by F Plasmids

T7 infection of host cells harboring an F plasmid is abortive, and T7 is, therefore, termed a "female-specific" phage (286). After initial results according to which two adjacent F-factor genes, *pifA* and *pifB*, determine an obstruction of late T7 mRNA translation, a number of papers described membrane permeability changes and a cessation of macromolecular syntheses ahead of late translation (for references, see Condit [78] and Hausmann [165]). Recently, Remes and Elseviers (384) proved that adenosine 5'-triphosphate leakage is not causally related to T7 exclusion. Despite numerous attempts to solve this problem, the ultimate cause of abortive development and the respective roles of chromosomal, plasmid, and phage genes in T7-infected F<sup>+</sup> cells are not completely clear. The F-plasmid genes responsible for T7 exclusion were recently cloned (465) and shown to be localized at the 38.3 position on the F-plasmid map (353).

Schell et al. (429) and Williams and Meynell (524) had reported an F-mediated exclusion of T3, whereas Hausmann (163) and Studier (490) failed to detect such a phenomenon. This discrepancy is explained by the fact that the phages used by the former authors were actually T7 strains (490). T3 can not be termed "female specific." It is to be hoped that a comparison of T3 and T7 will lead to the identification of the phage function(s) which confer resistance and sensitivity, respectively, to F<sup>+</sup> *pif* gene action.

### P1 Prophage

The impaired growth of T3 and T7 on P1-lysogenic cells has been known for a long time (122, 138, 268). Hausmann noted that gene expression declines sharply from the early to the late genes and that this drop occurs more rapidly in the case of T7 (163).

Krüger and co-workers (251; D. H. Krüger, S. Hansen, M. Reuter, and C. Schroeder, manuscript in preparation) have shown that the block on T3 and T7 development is caused by the restriction enzyme *EcoP1*. However, no cleavage of T3 or T7 DNA in P1-lysogenic cells is observed. It was proved that T3 (*sam*<sup>+</sup>), compared with T7 and T3*sam*<sup>-</sup> mutants, has a superior ability to express its genes and damage the cell.

Restriction endonuclease *EcoP1* molecules appear to bind to T3 and T7 DNAs and, similarly to repressors, inhibit transcription. DNA cleavage is prevented by the phage *ocr*<sup>+</sup> function. Due to the expression of SAMase and the resulting lack of SAM in the T3-infected cell, the repressor-like binding of the P1 endonuclease is less stable. T3 and T7 *ocr*<sup>-</sup> mutants are subject to in vivo DNA cleavage and methylation by restriction endonuclease *EcoP1*. The investigation of in vivo restriction of phages T3 and T7 by *EcoP1* has aided the dissection of the different levels on which the *sam*<sup>+</sup> and *ocr*<sup>+</sup> functions influence the interaction between DNA and restriction enzymes (see Interactions with the DNA Modification and Restriction System of the Host: Functions of Gene 0.3).

It is known that *EcoP1* and *EcoP15* show differences from the type I restriction enzymes in their in vitro properties (11, 151, 542) and, moreover, recognize a shorter nucleotide sequence (16, 152)—which should occur with a higher probability within a given DNA molecule—than does *EcoB* or *EcoK* (214, 264, 381, 468). It is still not clear which special feature of T3 and T7 is responsible for the strong repressor-like effect of the P1 restriction enzyme in vivo. In contrast to the drastic in vivo effect of *EcoP1* on T3 and T7 DNAs, phage lambda and the plasmid pSF2124 DNAs are restricted by *EcoP1* in vivo in the usual manner, which also obtains for the restriction of lambda and *ocr*<sup>-</sup> derivatives of T3 and T7 by *EcoB* and *EcoK* (no repressor effect) (11, 254, 387). In addition, *EcoP1*-mediated restriction of pSF2124 DNA can be completely overcome by *ocr*<sup>+</sup> activity (387).

### Antibiotic Resistance Plasmids and Colicinogenic Plasmids

Some older papers describe the inhibition of T3 or T7 phage growth by certain R plasmids (18, 312, 519). In none of these cases was the mechanism of exclusion (e.g., adsorption, abortive infection through incompatibility, DNA restriction) elucidated.

Our preliminary studies on the influence of the restriction endonucleases *EcoRI* and *EcoRII* on the propagation of T3 and T7 in vivo show that these restriction systems do not significantly inhibit virus growth (Krüger, unpublished data). This is not surprising in the case of *EcoRI*, since T3 and T7 DNAs have no recognition sites for *EcoRI* (17, 260, 405, 421). The DNA M/R system of plasmid R124, which has also been termed *EcoRIII* (464) and which corresponds to a type I restriction enzyme (S. G. Hughes, personal communication), does not restrict T3 phage, on account of the protective effect of the

*ocr*<sup>+</sup> gene function; T3*ocr*<sup>-</sup> mutants are restricted and modified in vivo (D. H. Krüger, S. Hansen, and L. S. Chernin, manuscript in preparation).

Various colicinogenic plasmids interfere with T3 or T7 growth not on the level of adsorption or DNA restriction, but in an abortive infection through incompatibility (146, 168).

### *Shigella sonnei* D<sub>2</sub> 371-48

The infection of *S. sonnei* D<sub>2</sub> 371-48 by T7 phage ends abortively. Though DNA synthesis begins as usual, it is followed by extensive phage DNA breakdown for which a suicidal (wild-type) phage function *ss*<sup>+</sup> is responsible, since T7*ss*<sup>-</sup> mutants are capable of normal growth in the *Shigella* strain. During infection of *E. coli* host cells, the *ss*<sup>+</sup> phage function is compensated by a "suicide inhibition function" of the cell (*sin*<sup>+</sup>), whereas *S. sonnei* D<sub>2</sub> is *sin* (cp. 164, 165). The *ss* marker was recently localized within the gene coding for the major head protein (gene 10) (513). The fact that a gene 10 mutation counteracts the suicidal behavior that normally occurs upon infection of *S. sonnei* D<sub>2</sub> (*sin*) suggests that an interaction between a host function (*sin*<sup>+</sup>) and gp10 of T7 may normally take place during phage head morphogenesis. The abortive infection of *S. sonnei* D<sub>2</sub> is caused by the block of phage assembly; the degradation of T7 DNA is a secondary effect (513). T3 growth is unperturbed on *S. sonnei* D<sub>2</sub> (203); T3 thus behaves as a "natural" *ss*<sup>-</sup> mutant. This is corroborated by the apparent nonhomologies between the gene 10 base sequences of T3 and T7 (89, 203).

Because of the facile discrimination between *ss*<sup>+</sup> and *ss*<sup>-</sup> T7 phage with the help of *S. sonnei* D<sub>2</sub>, these markers have been used as experimental tools for in vitro DNA recombination and packaging (294, 399, 413).

## SPECIAL VIRUS-VIRUS AND VIRUS-HOST INTERACTIONS

### T3 as a Facultative Temperate Phage

Lysogeny of bacterial cells has been studied in detail in the cases of such temperate phages as lambda and P22 (19, 118, 181, 500). The most important criteria for lysogenization are the perpetuation of prophage as part of the bacterial replicating system and the ability of the lysogenic cell to produce progeny phage (prophage induction) without reinfection. Phage-coded repressor molecules prevent the expression of lytic phage functions and are also responsible for the superinfection immunity of the cells to homologous phage. Due to the intracellular state of the phage, it is resistant to phage antiserum treatment. In addition to lysogeny, there are other

less intimate virus-cell interactions, called pseudolysogeny or carrier state, which have not been strictly defined (19, 125, 174, 209, 278, 505, 518).

In the 1950s Fraser and co-workers (124, 125, 182) noticed that after infection by phage T3 of *E. coli* cells starved in buffer, there is no immediate phage replication and special so-called late-lysing complexes are formed. In 1972 Krüger et al. (252) observed that upon infection of starved *E. coli* B cells, a fundamental difference between T3 and T7 becomes evident. Whereas, T7 induces a normal lytic cycle, the T3 genome forms a complex with the cell. During slow cell growth in minimal medium the phage genome is coreplicated, and the daughter cells carry the viral information. A transfer of T3-lysogenic cells into rich medium causes induction of T3 prophage, and a complete lytic cycle is performed. The facultative temperate behavior is connected to the T3 SAMase (*sam*<sup>+</sup>) gene (243). Apparently the T3 genome, which is nonmethylated due to SAMase action during the preceding growth cycle (see Interactions with the DNA Modification and Restriction System of the Host: Functions of Gene 0.3), cannot be expressed in starved cells. In contrast, SAMase-negative phage with methylated genomes (T7, T3*sam*<sup>-</sup>) are expressed in starved host cells. When, however, T3*sam*<sup>-</sup> phage are propagated under conditions where DNA methylation is prevented by preinfection with SAMase-proficient phage, these unmethylated *sam*<sup>-</sup> phage also establish a temperate interaction upon subsequent infection of starved cells (243). It is still a matter of speculation why the degree of methylation is so important for the behavior of T3 in starved host cells but not in normal cells. During the lysogenic state, not even the leftmost portion of the T3 genome is expressed (D. H. Krüger and C. Schroeder, unpublished data). Replication of the T3 prophage is thus under exclusive control of the *E. coli* replication system. Incubation of T3-lysogenic cells in minimal medium plus T3 antiserum has no influence on the lysogenic state; i.e., reinfection (carrier state) cannot be the cause for T3 perpetuation.

In contrast to authentic lysogeny, e.g., by phage lambda, T3 lysogeny is unstable and not maintained by phage-coded repressor action. The absence of a phage repressor is also manifested in the lack of superinfection immunity (Krüger, unpublished data). Up to now it is not clear whether T3 prophage is incorporated in the *E. coli* chromosome or whether it is "inherited" as a plasmid.

Thus, under special circumstances the normally virulent phage T3 can obey criteria of temperance. We prefer to call this type of virus-host interaction *lysogeny* to emphasize the abil-

ity of the growing cell to lyse after induction of provirus. A decision for a certain term would demand a more detailed knowledge of the underlying molecular processes. Possibly, *intracellular persistence* would be more appropriate, since the term *lysogeny* is traditionally associated with high stability and the expression of at least one phage gene product, the repressor protein. In its dependence on the phage and possibly the methylation degree of its genome, on the one hand, and the physiological state of the host cell, on the other, temporary T3 lysogeny provides an interesting system for studying the regulation of host "rescue" gene expression programs (e.g., the stringent control pattern [130, 329] or the SOS pattern [527]) on the level of transcriptional control and their possible relation to DNA methylation. First indications of a link between differentiation processes and DNA methylation have also been reported for eucaryotic systems (4, 75, 308, 501, 547).

### *E. coli* transfection

The utilization of transfection studies on *Enterobacteriaceae* for the solution of different problems of molecular biology was recently reviewed by Benzinger (28). Progress in the transfection of *E. coli* spheroplasts by T7 DNA has been made by the groups of Benzinger (29, 176, 265) and Wackernagel (104, 448, 517). Ehrlich et al. (119) studied the transfection of  $\text{CaCl}_2$ -treated *E. coli* by T7 DNA.

T7 DNA is infective in the double-stranded and the single-stranded state (265, 297, 448, 517). Transfection by T7 single-stranded DNA was observed to depend on a functional DNA polymerase III in *polAB* cells, whereas transfection with native T7 DNA was independent of host DNA polymerase (448). The infectivity of double-stranded DNA in vivo is significantly reduced by the *recBC* nuclease, probably by an attack on the free ends of the linear duplex (29, 119, 448, 517). An in vitro erosion of the 5' ends of these duplexes, i.e., the creation 3'-terminal single strands, does not lower, but rather enhances, infectivity (104, 119, 517), since these are poorer substrates for the *recBC* enzyme, and, furthermore, circularization of "terminally eroded" DNA takes place within the cell, making it completely insensitive to *recBC* digestion (104, 448, 517). However, when both strands of the terminal redundancies have been degraded in vitro, the DNA loses its infectivity. This is yet another proof of the essentiality of the terminal redundancies for DNA replication and maturation (104).

Transfection efficiency of restriction-proficient cells by T7 DNA is substantially reduced

compared with restriction-deficient cells, which is explained by restriction of the entering DNA molecule (119). The fact that the *E. coli* strain used for propagating the phage from which the DNA was extracted had no influence on the degree of restriction (119) is explained by the fact that *ocr*<sup>+</sup> action prevented DNA modification (245). During normal infection, incoming DNA is protected against restriction by the *ocr*<sup>+</sup> function (see Interactions with the DNA Modification and Restriction System of the Host: Functions of Gene 0.3), but in the transfected cell this phage function is, of course, expressed too late. Interestingly, the 1,000-fold reduction of transfection efficiency of T7 DNA by restriction endonuclease *EcoK* (119) corresponds exactly to the drop in efficiency of plating of T7*ocr*<sup>-</sup> mutants when they are plated on *E. coli* K (254).

Analogous to the fact that the DNA restriction system of the host does not inhibit T7 growth upon normal infection but upon transfection, the *recBC* nuclease seems only to interfere with transfection, and not with normal T7 infection (517). Phage whose DNA goes through a *recBC*-sensitive stage in their life cycle have evolved protective measures against the enzyme (28). This protection has been most thoroughly elucidated in the case of the gamma protein of phage lambda (216, 420). Sakaki (419) found that besides lambda and other phages, T3 and T7 are also capable of inactivating the *recBC* nuclease and that inactivation depends on phage gene expression. Wackernagel and Hermanns (516) came to the conclusion that the anti-*recBC* effect is caused by a late T7 protein. This protein was isolated and partially characterized (349). Unfortunately, no mutants in the corresponding gene are known.

### Problem of Heterologous and Homologous Superinfection Exclusion

Hausmann et al. (171, 173) described the effect that in homologous (T3  $\times$  T3 and T7  $\times$  T7) crosses, the phage are compatible and both parental genomes are replicated and expressed, but heterologous mixed infection by T3 plus T7 results in mutual exclusion and only 5% of the mixed-infected cells produce both phage species. Beier (Ph.D. thesis) found that upon simultaneous infection, exclusion only takes place in the heterologous system, whereas upon superinfection, such an exclusion was postulated to occur in heterologous as well as homologous infection. The marker determining exclusion appears to reside in gene 1 (phage RNA polymerase) (Beier, Ph.D. thesis), indicating a role of the different template specificities of the phage RNA polym-



erases. Superinfection experiments with T3- and T7-preinfected cells confirmed the role of gene 1 (299), and the results of McAllister and Barrett (Table 1 of reference 299) let us conclude that gene 0.3 cannot be involved to any extent. Benbasat et al. (27) claim that after T7 infection, all superinfecting phage are excluded, since they adsorb but do not inject their DNA; this effect is supposed to be completely reversible by chloramphenicol; i.e., it is supposed to require protein synthesis by the first infecting phage. Hirsch-Kauffmann et al. (193) express the most extreme opinion. According to their results, T3 and T7 exclude each other in homologous as well as heterologous co- and superinfections. Only one phage genome should be able to enter the cell and block the entry of any other phage in a step subsequent to adsorption. This exclusion is allegedly mediated by gene 0.3 (called "M" gene by these authors [see Other Functions Ascribed to Gene 0.3]), without a requirement for RNA or protein synthesis (no chloramphenicol effect) directed by the excluding virus. Due to the postulated exclusion, no ultraviolet multiplicity reactivation of T3 and T7 should occur (193); however, multiplicity reactivation has been observed by other authors (45, 46).

As contradictory as these data are (Exclusion only on superinfection or also in coinfection? Exclusion only between heterologous or also between homologous phage? Role of gene 0.3 or gene 1 for exclusion? Necessity of gene expression of the first phage for the exclusion of all following phage?), they are as a whole opposed to quite self-evident facts of the molecular biology of T3 and T7. It is a well-known fact that amber mutants (23, 166, 483) and temperature-sensitive mutants (253, 442) efficiently complement each other in coinfecting cells and that complementation is also observed in heterologous mixed infections (168). Obviously, complementation depends on the expression of at least two different phage genomes in the cell. The same is true for recombination experiments. Recombination frequencies in homologous crosses are normal (23, 483, 485, 494, 508). The recombination frequencies in heterologous T3  $\times$  T7 crosses are indeed lower than in homologous crosses (171; Beier, Ph.D. thesis), but the reason for this is that recombinants resulting from one (or any uneven number of) recombinational event are not viable and therefore not detectable, because recombinant DNA molecules carrying nonidentical terminal "redundancies" cannot form concatemers (22, 89, 104). (During experiments leading to the establishment of recombinational maps of T3 and T7, reproducibility is improved if phage infection is synchronized by

applying KCN. This measure has no influence on the expression of gene 0.3 or 1.)

Bräutigam and Sauerbier (40) found a gene dosage effect depending on the multiplicity of infection of ultraviolet-irradiated T7 gene 1 mutants. Other authors found that net T3 DNA synthesis increases with the multiplicity of infection by T3 wild type (W. Mann, H. Musielski, I. Scheiber, and I. Laue, Abstr. Commun. Annu. Meet. Biochem. Soc. GDR, 11th, Halle/Saale, German Democratic Republic, 1979, B3). In connection with other experiments, Krüger and co-workers have shown that, for example, after preinfection of *E. coli* with T3 gene 4 or gene 6 amber mutants, subsequently infecting T3 $_{sam}^-$  normally replicates (243) and that preinfection by ultraviolet-irradiated T3 or T7 phage allows normal growth of T3/R7 and T7/D111 (245) and T3 and T7 wild types (Krüger, unpublished data). In all these cases a normal expression of gene 0.3 (and in the case of the amber mutants, also of gene 1) of the preinfecting phage takes place. All these results, including the complementation and recombination mapping data, imply that homologous or heterologous co- or superinfection immunity does not exist.

Since all experiments relating to this matter have been performed with phage mutants or ultraviolet-irradiated phage, there is, of course, the possibility that some kind of exclusion exists between wild-type phage. If there is a certain degree of mutual exclusion in the case of heterologous T3  $\times$  T7 crosses (171, 173), then it is certainly not related to gene 0.3 function.

The arguments given above hold only for simultaneous infection or for superinfection involving a delay of maximally 5 min (37°C). It is evident that a secondary infection after a longer lapse of time is bound to be unsuccessful because the processes in the infected cell are out of control.

### Interactions with Eucaryotic Systems

Prompted by the spectacular finding of Merrill and co-workers (134, 311) that  $\lambda$ gal induces galactose-1-phosphate uridylyltransferase in galactosemic human cells and by successful phage transduction of *E. coli* genes into plant cells (101), the interactions between phage and eucaryotic systems attracted the interest of molecular biologists (cp. 309, 310).

Up to now, neither complete bacteriophage reproduction nor stable integration of whole phage genomes in higher eucaryotic cells has been described. However, the induction of plant tumors by the Ti plasmid of *Agrobacterium tumefaciens* represents a natural case of stable

incorporation of bacterial genes in eucaryotic cells (273, 428, 540).

After infection of *Hordeum vulgare* protoplasts by T3, the phage genome is expressed to a limited extent (54). Regarding the fate of T7 DNA after experimental introduction into animal cells, there are conflicting results: Leavitt et al. (266) demonstrated the uptake of T7 DNA into the nucleus of Syrian hamster embryonic cells and its persistence throughout several cell division cycles; Kao et al. (215), in contrast, observed rapid digestion of T7 DNA after its entry into human skin fibroblasts. These and other results (see reference 139) invite the speculation that, analogous to the situation in bacteria (11, 37, 504), restriction systems with the ability of degrading foreign DNA also exist in eucaryotic cells (47, 416). Maybe these restriction systems are not yet fully expressed in embryonic, compared with adult, cells.

Richter et al. (389) detected a limited transcription and translation of T3 and T7 DNA in a cell-free system of yeast mitochondria. That phage DNA expression is not confined to the procaryote-like mitochondrial system was demonstrated by Anderson et al. (9), who found that T3 and T7 early mRNA's are translated on the 80S ribosomes of a mammalian cell-free cytoplasmic system. The translation efficiency of gene 0.3 mRNA is higher when using RNase III-processed mRNA than with polycistronic RNA. Capping of mRNA (modification of the 5' end which facilitates translation of eucaryotic mRNA) seems not to be necessary for phage mRNA translation (9).

The purpose of these studies is twofold: first, they may help to illuminate the differences and correspondences in the molecular biology of pro- and eucaryotes, and, second, these studies could be of practical value for future experimental gene transfer with recombinant DNA from pro- to eucaryotes.

#### ACKNOWLEDGMENTS

We gratefully acknowledge the unfailing support of our work extended to us by H. A. Rosenthal. Thanks are due to many colleagues for personal communications, preprints and reprints of their papers, and for the gift of bacterial and phage strains: W. Arber, Hildburg Beier, Ulla Günthert, R. Hausmann, J. Hofemeister, S. G. Hughes, S. Michel, F. W. Studier, and W. Wackernagel. We also thank T. G. Chanishvili, L. S. Chernin, Liana Gachechiladze, Sigrud Hansen, W. Presber, Monika Reuter, and G. B. Zavlzelsky for dedicated collaboration in our T3 and T7 research.

#### LITERATURE CITED

- Abelson, J. 1979. RNA processing and the intervening sequence problem. *Annu. Rev. Biochem.* **48**:1035-1069.
- Abraham, K. A., K. J. Andersen, and A. Rognes. 1972. Studies on DNA-dependent RNA polymerase from *E. coli*. Variations of the enzyme activity during growth. *Biochem. J.* **129**:291-299.
- Adams, M. H., and E. Wade. 1954. Classification of bacterial viruses: the relationship of two Serratia phages to coli-dysentery phages T3, T7, and D44. *J. Bacteriol.* **68**:320-325.
- Adams, R. L. P. 1973. Delayed methylation of DNA in developing sea urchin embryos. *Nature (London)* **244**:27-29.
- Adhya, S., and M. Gottesman. 1978. Control of transcription termination. *Annu. Rev. Biochem.* **47**:967-996.
- Adhya, S., P. Sarkar, D. Valenzuela, and U. Maitra. 1979. Termination of transcription by *E. coli* RNA polymerase: influence of secondary structure of RNA transcripts on rho-independent and rho-dependent termination. *Proc. Natl. Acad. Sci. U.S.A.* **76**:1613-1617.
- Adler, S., and P. Modrich. 1979. T7-induced DNA polymerase. Characterization of associated exonuclease activities and resolution into biologically active subunits. *J. Biol. Chem.* **254**:11605-11614.
- Alberts, B. M., and R. Sternglanz. 1977. Recent excitement in the DNA replication problem. *Nature (London)* **269**:655-661.
- Anderson, C. W., J. R. Atkins, and J. J. Dunn. 1976. Bacteriophage T3 and T7 early RNAs are translated by eucaryotic 80S ribosomes: active phage T3 coded S-adenosylmethionine cleaving enzyme is synthesized. *Proc. Natl. Acad. Sci. U.S.A.* **73**:2752-2756.
- Arai, K.-I., and A. Kornberg. 1979. A general priming system employing only *dnaB* protein and primase for DNA replication. *Proc. Natl. Acad. Sci. U.S.A.* **76**:4308-4312.
- Arber, W. 1974. DNA modification and restriction. *Prog. Nucleic Acid Res. Mol. Biol.* **14**:1-37.
- Arber, W. 1977. Open question: What is the function of DNA restriction enzymes. *Trends Biochem. Sci.* **2**:N175-N176.
- Arber, W. 1979. Promotion and limitation of genetic exchange. *Science* **205**:361-365.
- Arber, W., R. Yuan, and T. A. Bickle. 1975. Strain-specific modification and restriction of DNA in bacteria. *FEBS (Fed. Eur. Biochem. Soc.) Proc. Meet.* **34**:3-22.
- Auer, B., U. Günthert, E. F. Wagner, and M. Schweiger. 1979. Is the DNA of virus T7 methylated? *J. Gen. Virol.* **44**:609-613.
- Bächi, B., J. Reiser, and V. Pirotta. 1979. Methylation and cleavage sequences of the *EcoPI* restriction-modification enzyme. *J. Mol. Biol.* **128**:143-163.
- Bailey, J. N., D. R. Dembinski, and W. T. McAllister. 1980. Derivation of a restriction map of bacteriophage T3 DNA and comparison with the map of bacteriophage T7 DNA. *J. Virol.* **35**:176-183.
- Bannister, D., and S. W. Glover. 1968. Restriction and modification of bacteriophages by R<sup>+</sup>

- strains of *E. coli* K12. Biochem. Biophys. Res. Commun. 30:735-738.
19. Barksdale, L., and S. B. Arden. 1974. Persisting bacteriophage infections, lysogeny, and phage conservation. Annu. Rev. Microbiol. 28: 265-299.
  20. Bautz, E. K. F. 1973. Initiation of transcription by RNA polymerases of *E. coli* and phage T3. FEBS Lett. 36:123-129.
  21. Becker, A., G. Lyn, M. Geftter, and J. Hurwitz. 1967. The enzymatic repair of DNA. II. Characterization of phage-induced sealase. Proc. Natl. Acad. Sci. U.S.A. 58:1996-2003.
  22. Beier, H., M. Golomb, and M. Chamberlin. 1977. Isolation of recombinants between T7 and T3 bacteriophages and their use in vitro transcriptional mapping. J. Virol. 21:753-765.
  23. Beier, H., and R. Hausmann. 1973. Genetic map of bacteriophage T3. J. Virol. 12:417-419.
  24. Beier, H., and R. Hausmann. 1974. T3  $\times$  T7 crosses leading to recombinant RNA polymerases. Nature (London) 251:538-540.
  25. Belogurov, A. A., and G. B. Zavilgelsky. 1977. Significance of the *EcoK* restriction system for DNA cross-link repair. Dokl. Akad. Nauk SSSR 235:208-211. (In Russian.)
  26. Belogurov, A. A., and G. B. Zavilgelsky. 1979. Repair of interstrand cross links in DNA: Role of the *E. coli* K restriction system. Mol. Biol. (USSR) 13:160-168. (In Russian.)
  27. Benbasat, I. A., K. B. Burck, and R. C. Miller. 1978. Superinfection exclusion and lack of conservative transfer of bacteriophage T7 DNA. Virology 87:164-171.
  28. Benzinger, R. 1978. Transfection of *Enterobacteriaceae* and its applications. Microbiol. Rev. 42:194-236.
  29. Benzinger, R., L. W. Enquist, and A. Skalka. 1975. Transfection of *Escherichia coli* spheroplasts. V. Activity of *recBC* nuclease in *rec<sup>+</sup>* and *rec<sup>-</sup>* spheroplasts measured with different forms of bacteriophage DNA. J. Virol. 15:861-871.
  30. Bernstein, C., and H. Bernstein. 1974. Coiled rings of DNA released from cells infected with bacteriophages T7 or T4 or from uninfected *Escherichia coli*. J. Virol. 13:1346-1355.
  31. Bick, M. D. 1974. Electron microscopic analysis of T7 DNA synthesized in vitro. J. Biol. Chem. 249:2974-2984.
  32. Bishayee, S., A. Yudelevich, and U. Maitra. 1976. Specificity of chain initiation by bacteriophage T7-induced RNA polymerase. Biochem. Biophys. Res. Commun. 68:626-633.
  33. Bogdanova, E. S., Z. M. Gorlenko, and E. M. Khourgess. 1974. On the effect of bacteriophage infection on host RNA polymerase. Mol. Gen. Genet. 133:261-272.
  34. Boothroyd, J. C., and R. S. Hayward. 1979. New genes and promoters suggested by the DNA sequence near the end of the coliphage T7 early operon. Nucleic Acids Res. 7:1931-1943.
  35. Bordier, C. 1974. Inhibition of rifampicin-resistant RNA synthesis by rifampicin-RNA polymerase complexes. FEBS Lett. 45:259-262.
  36. Bordier, C., and J. Dubochet. 1974. Electron microscopic localization of the binding sites of *E. coli* RNA polymerase in the early promoter region of T7 DNA. Eur. J. Biochem. 44:617-624.
  37. Boyer, H. W. 1971. DNA restriction and modification mechanisms in bacteria. Annu. Rev. Microbiol. 25:153-176.
  38. Braun, V., and K. Hantke. 1974. Biochemistry of bacterial cell envelopes. Annu. Rev. Biochem. 43:89-121.
  39. Bräutigam, A. R., and W. Sauerbier. 1973. Transcription unit mapping in bacteriophage T7. I. In vivo transcription by *Escherichia coli* RNA polymerase. J. Virol. 12:882-886.
  40. Bräutigam, A. R., and W. Sauerbier. 1974. Transcription unit mapping in bacteriophage T7. II. Proportionality of number of gene copies, mRNA, and gene product. J. Virol. 13: 1110-1117.
  41. Brown, P. D., C. L. Peebles, and N. R. Cozzarelli. 1979. A topoisomerase from *E. coli* related to DNA gyrase. Proc. Natl. Acad. Sci. U.S.A. 12:6110-6114.
  42. Brunel, F., and J. Davison. 1979. Restriction insensitivity in bacteriophage T5. III. Characterization of *EcoRI*-sensitive mutants by restriction analysis. J. Mol. Biol. 128:527-543.
  43. Brunovskis, I., and R. D. Burns. 1973. Growth of coliphage T7 in *Salmonella typhimurium*. J. Virol. 11:621-629.
  44. Brunovskis, I., and W. C. Summers. 1972. The process of infection with coliphage T7. VI. A phage gene controlling shutoff of host RNA synthesis. Virology 50:322-327.
  45. Burck, K. B., and R. C. Miller. 1978. Marker rescue and partial replication of bacteriophage T7 DNA. Proc. Natl. Acad. Sci. U.S.A. 12: 6144-6148.
  46. Burck, K. B., D. G. Scraba, and R. C. Miller. 1979. Electron microscopic analysis of partially replicated bacteriophage T7 DNA. J. Virol. 32: 606-613.
  47. Burdon, R. H., A. C. B. Cato, and I. S. Villadsen. 1979. S-Adenosyl-L-methionine-dependent deoxyribonucleases of eukaryotic origin. FEBS (Fed. Eur. Biochem. Soc.) Proc. Meet. 51:159-168.
  48. Burton, W. G., C. T. Grabowy, and R. Sager. 1979. Role of methylation in the modification and restriction of chloroplast DNA in *Chlamydomonas*. Proc. Natl. Acad. Sci. U.S.A. 76: 1390-1394.
  49. Cairns, J., G. S. Stent, and J. D. Watson. (ed.). 1966. Phage and the origins of molecular biology. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
  50. Campbell, J. L., C. C. Richardson, and F. W. Studier. 1978. Genetic recombination and complementation between bacteriophage T7 and cloned fragments of T7 DNA. Proc. Natl. Acad. Sci. U.S.A. 75:2276-2280.
  51. Campbell, J. L., F. Tamanoi, C. C. Richard-

- son, and F. W. Studier. 1979. Cloning of the T7 genome in *E. coli*: use of recombination between cloned sequences and bacteriophage T7 to identify genes involved in recombination and a clone containing the origin of T7 DNA replication. Cold Spring Harbor Symp. Quant. Biol. 43:441-448.
52. Cantoni, G. L. 1975. Biological methylation: selected aspects. Annu. Rev. Biochem. 44:435-451.
53. Carlson, K. 1968. Intracellular fate of DNA from T7 bacteriophage. J. Virol. 2:1230-1233.
54. Carlson, P. S. 1973. The use of protoplasts for genetic research. Proc. Natl. Acad. Sci. U.S.A. 70:598-602.
55. Casjens, S., and J. King. 1975. Virus assembly. Annu. Rev. Biochem. 44:555-611.
56. Cech, C. L., and W. R. McClure. 1980. Characterization of RNA polymerase-T7 promoter binary complexes. Biochemistry 19:2440-2454.
57. Center, M. S. 1972. Bacteriophage T7-induced endonuclease II: purification and properties of the enzyme. J. Biol. Chem. 247:146-156.
58. Center, M. S. 1972. Replicative intermediates of bacteriophage T7 DNA. J. Virol. 10:115-123.
59. Center, M. S. 1973. Bacteriophage T7 DNA synthesis in isolated DNA membrane complexes. J. Virol. 12:847-854.
60. Center, M. S. 1975. Role of gene 2 in bacteriophage T7 DNA synthesis. J. Virol. 16:94-100.
61. Center, M. S., and C. C. Richardson. 1970. An endonuclease induced after infection of *E. coli* with bacteriophage T7. I. Purification and properties of the enzyme. J. Biol. Chem. 245:6285-6291.
62. Chakrabarty, S. L., and L. Gorini. 1975. A link between streptomycin and rifampicin mutation. Proc. Natl. Acad. Sci. U.S.A. 72:2084-2087.
63. Chakrabarty, S. L., and L. Gorini. 1977. Interaction between mutations of ribosomes and RNA polymerase: a pair of *strA* and *rif* mutants individually temperature-insensitive but temperature-sensitive in combination. Proc. Natl. Acad. Sci. U.S.A. 74:1157-1161.
64. Chakraborty, P. R., P. Bandyopadhyay, P. Huang, and U. Maitra. 1974. Fidelity of in vitro transcription of T3 DNA by bacteriophage T3-induced RNA polymerase and by *E. coli* RNA polymerase. J. Biol. Chem. 249:6901-6909.
65. Chakraborty, P. R., R. A. Salvo, H. K. Majumder, and U. Maitra. 1977. Further characterization of bacteriophage T3-induced RNA polymerase. J. Biol. Chem. 252:6485-6493.
66. Chakraborty, P. R., P. Sarkar, H. H. Huang, and U. Maitra. 1973. Studies on T3-induced RNA polymerase from bacteriophage T3-infected *E. coli* cells. J. Biol. Chem. 248:6637-6646.
67. Chamberlin, M. 1974. Isolation and characterization of prototrophic mutants of *Escherichia coli* unable to support the intracellular growth of T7. J. Virol. 14:509-516.
68. Chamberlin, M. 1976. Interaction of RNA polymerase with the RNA template, p. 159-191. In R. Losick and M. Chamberlin (ed.), RNA polymerase. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
69. Chamberlin, M. J. 1974. The selectivity of transcription. Annu. Rev. Biochem. 43:721-775.
70. Chamberlin, M. J., J. McGrath, and L. Waskell. 1970. New RNA polymerase from *E. coli* infected with bacteriophage T7. Nature (London) 228:227-231.
71. Chamberlin, M. J., and J. Ring. 1973. Characterization of T7-specific RNA polymerase. I. General properties of the enzymatic reaction and the template specificity of the enzyme. J. Biol. Chem. 248:2235-2244.
72. Champoux, J. J. 1978. Proteins that affect DNA conformation. Annu. Rev. Biochem. 47:449-479.
73. Chao, L., and J. F. Speyer. 1973. A new form of RNA polymerase isolated from *E. coli*. Biochem. Biophys. Res. Commun. 51:399-405.
74. Cherny, D. I., A. A. Aleksandrov, M. I. Zardnaya, Y. N. Kosaganov, and Y. S. Lazurkin. 1977. Investigation of the binding of *E. coli* RNA polymerase to DNA from bacteriophages T2 and T7 by kinetic formaldehyde method and electron microscopy. Eur. J. Biochem. 79:309-317.
75. Christman, J. K., P. Price, L. Pedrinan, and G. Acs. 1977. Correlation between hypomethylation of DNA and expression of globin genes in Friend erythroleukemia cells. Eur. J. Biochem. 81:53-61.
76. Clark, R. W., G. H. Wever, and J. S. Wiberg. 1980. High-molecular weight DNA and the sedimentation coefficient: a new perspective based on DNA from T7 bacteriophage and two novel forms of T4 bacteriophage. J. Virol. 33:438-448.
77. Coleman, J. E. 1974. The role of Zn(II) in transcription by T7 RNA polymerase. Biochem. Biophys. Res. Commun. 60:641-648.
78. Condit, R. 1976. Who killed T7? Nature (London) 260:287-288.
79. Condit, R. C. 1975. F factor-mediated inhibition of bacteriophage T7 growth: increased membrane permeability and decreased ATP levels following T7 infection of male *E. coli*. J. Mol. Biol. 98:45-56.
80. Conley, R. R., and V. Pigiet. 1978. In vivo distribution of phosphothioredoxin in *E. coli*. J. Biol. Chem. 253:5568-5572.
81. Coulondre, C., J. H. Miller, P. J. Farabaugh, and W. Gilbert. 1978. Molecular basis of base substitution hotspots in *E. coli*. Nature (London) 274:775-780.
82. Cronan, J. E., Jr., R. Reed, F. R. Taylor, and M. B. Jackson. 1979. Properties and biosynthesis of cyclopropane fatty acids in *Escherichia coli*. J. Bacteriol. 138:118-121.
83. Darlix, J.-L. 1973. The functions of rho in T7-DNA transcription in vitro. Eur. J. Biochem. 35:517-526.
84. Darlix, J.-L. 1974. Rho, a factor causing the modulation of early T7 genes transcription. Biochimie 56:693-701.
85. Darlix, J. L., and J.-P. Dausse. 1975. Locali-

- zation of *E. coli* RNA polymerase initiation sites in T7 DNA early promoter region. *FEBS Lett.* **50**:214-218.
86. Darlix, J.-L., and M. Horaist. 1975. Existence and possible roles of transcriptional barriers in T7 DNA early region as shown by electron microscopy. *Nature (London)* **256**:288-292.
  87. Dausse, J.-P., A. Sentenac, and P. Fromageot. 1975. Interaction of RNA polymerase from *E. coli* with DNA. Analysis of T7 DNA early promoter sites. *Eur. J. Biochem.* **57**:569-578.
  88. Davis, R. W., and R. W. Hyman. 1970. Physical locations of the in vitro RNA initiation site and termination sites of T7M DNA. Cold Spring Harbor Symp. Quant. Biol. **35**:269-281.
  89. Davis, R. W., and R. W. Hyman. 1971. A study in evolution: the DNA base sequence homology between coliphages T7 and T3. *J. Mol. Biol.* **62**:287-301.
  90. Davison, J., and F. Brunel. 1979. Restriction insensitivity in bacteriophage T5. I. Genetic characterization of mutants sensitive to *EcoRI* restriction. *J. Virol.* **29**:11-16.
  91. Davison, J., and F. Brunel. 1979. Restriction insensitivity in bacteriophage T5. II. Lack of *EcoRI* modification in T5<sup>+</sup> and T5<sup>ris</sup> mutants. *J. Virol.* **29**:17-20.
  92. Day, R. S., III. 1977. UV-induced alleviation of K-specific restriction of bacteriophage  $\lambda$ . *J. Virol.* **21**:1249-1251.
  93. DeMartini, M., S. Halegona, and M. Inouye. 1975. Lysozymes from bacteriophages T3 and T7. *J. Virol.* **16**:459-461.
  94. Demerec, M., and U. Fano. 1945. Bacteriophage-resistant mutants in *E. coli*. *Genetics* **30**:119-136.
  95. Denhardt, D. T. 1979. DNA gyrase and DNA unwinding. *Nature (London)* **280**:196-198.
  96. DeWynngaert, M. A., and D. C. Hinkle. 1979. Bacterial mutants affecting phage T7 DNA replication produce RNA polymerase resistant to inhibition by the T7 gene 2 protein. *J. Biol. Chem.* **254**:11247-11253.
  97. DeWynngaert, M. A., and D. C. Hinkle. 1979. Involvement of DNA gyrase in replication and transcription of bacteriophage T7 DNA. *J. Virol.* **29**:529-535.
  98. DeWynngaert, M. A., and D. C. Hinkle. 1980. Characterization of the defects in bacteriophage T7 DNA synthesis during growth in the *Escherichia coli* mutant *tsnB*. *J. Virol.* **33**:780-788.
  99. Dharmalingam, K., and E. B. Goldberg. 1976. Phage-coded protein prevents restriction of unmodified progeny T4 DNA. *Nature (London)* **260**:454-456.
  100. Dharmgrongartama, B., S. P. Mahadik, and P. R. Srinivasan. 1973. Modification of RNA polymerase after T3 phage infection of *E. coli* B. *Proc. Natl. Acad. Sci. U.S.A.* **70**:2845-2849.
  101. Doy, C. H., P. M. Gresshoff, and B. G. Rolfe. 1973. Biological and molecular evidence for the transgenesis of genes from bacteria to plant cells. *Proc. Natl. Acad. Sci. U.S.A.* **70**:723-726.
  102. Dreiseikelmann, B., R. Eichenlaub, and W. Wackernagel. 1979. The effect of differential methylation by *E. coli* of plasmid DNA and phage T7 and lambda DNA in the cleavage by restriction endonuclease *MboI* from *Moraxella bovis*. *Biochim. Biophys. Acta* **562**:418-428.
  103. Dreiseikelmann, B., U. Steger, and W. Wackernagel. 1980. Length determination of the terminal redundant regions in the DNA of phage T7. *Mol. Gen. Genet.* **178**:237-240.
  104. Dreiseikelmann, B., and W. Wackernagel. 1978. The terminal redundant regions of bacteriophage T7 DNA. Their necessity for phage production studied by the infectivity of T7 DNA after modification by various exonucleases. *Mol. Gen. Genet.* **159**:321-328.
  105. Dressler, D. 1975. The recent excitement in the DNA growing point problem. *Annu. Rev. Microbiol.* **29**:525-559.
  106. Dressler, D., J. Wolfson, and M. Magazin. 1972. Initiation and reinitiation of DNA synthesis during replication of bacteriophage T7. *Proc. Natl. Acad. Sci. U.S.A.* **69**:998-1002.
  107. Dunn, J. J. 1976. RNaseIII cleavage of single-stranded RNA. Effect of ionic strength on the fidelity of cleavage. *J. Biol. Chem.* **251**:3807-3814.
  108. Dunn, J. J., C. W. Anderson, J. F. Atkins, D. C. Bartelt, and W. C. Crockett. 1976. Bacteriophage T7 and T3 as model systems for RNA synthesis and processing. *Prog. Nucleic Acid Res. Mol. Biol.* **19**:263-273.
  109. Dunn, J. J., F. A. Bautz, and E. K. F. Bautz. 1971. Different template specificities of phage T3 and T7 RNA polymerases. *Nature (London)* **230**:94-96.
  110. Dunn, J. J., E. Buzash-Pollert, and F. W. Studier. 1978. Mutations of bacteriophage T7 that affect initiation of synthesis of the gene 0.3 protein. *Proc. Natl. Acad. Sci. U.S.A.* **75**:2741-2745.
  111. Dunn, J. J., W. T. McAllister, and E. K. F. Bautz. 1972. Transcription in vitro of T3 DNA by *E. coli* and T3 RNA polymerases: analysis of the products in cell-free protein-synthesizing system. *Eur. J. Biochem.* **29**:500-508.
  112. Dunn, J. J., and F. W. Studier. 1973. T7 early RNAs are generated by site-specific cleavages. *Proc. Natl. Acad. Sci. U.S.A.* **70**:1559-1563.
  113. Dunn, J. J., and F. W. Studier. 1973. T7 early RNAs and *E. coli* rRNAs are cut from large precursor RNAs in vivo by RNaseIII. *Proc. Natl. Acad. Sci. U.S.A.* **70**:3296-3300.
  114. Dunn, J. J., and F. W. Studier. 1974. Processing, transcription, and translation of bacteriophage T7 mRNAs. *Brookhaven Symp. Biol.* **26**:267-276.
  115. Dunn, J. J., and F. W. Studier. 1975. Effect of RNaseIII cleavage on translation of bacteriophage T7 mRNAs. *J. Mol. Biol.* **99**:487-499.
  116. Dunn, J. J., and F. W. Studier. 1980. The transcription termination site at the end of the early region of bacteriophage T7 DNA. *Nucleic Acids Res.* **8**:2119-2132.
  117. Earnshaw, W. C., and S. R. Casjens. 1980. DNA packaging by the double-stranded bacteriophages. *Cell* **21**:319-331.

118. Echols, H. 1972. Developmental pathways for the temperate phage: lysis vs. lysogeny. *Annu. Rev. Genet.* **6**:157-190.
119. Ehrlich, S. D., V. Sgarbetta, and J. Lederberg. 1976. Transfection of restrictionless *E. coli* by bacteriophage T7 DNA: effect of in vitro erosion of DNA by lambda exonuclease. *J. Mol. Biol.* **105**:603-609.
120. Ennis, H. L., and K. D. Kievitt. 1977. Bacteriophage proteins associated with the bacterial membrane after bacteriophage T7 infection. *J. Virol.* **22**:561-567.
121. Eskin, B., J. A. Lautenberger, and S. Linn. 1973. Host-controlled modification and restriction of bacteriophage T7 by *Escherichia coli* B. *J. Virol.* **11**:1020-1023.
122. Eskridge, R. W., H. Weinfeld, and K. Paigen. 1967. Susceptibility of different coliphage genomes to host-controlled variation. *J. Bacteriol.* **93**:835-844.
123. Fenner, F., B. R. McAuslan, C. A. Mims, J. Sambrook, and D. O. White. 1974. The biology of animal viruses, 2nd ed. Academic Press, Inc., New York.
124. Fraser, D., and R. Dulbecco. 1953. A genetic analysis of the factors controlling the *ts* character in bacteriophage T3. Cold Spring Harbor Symp. Quant. Biol. **18**:15-17.
125. Fraser, D. K. 1957. Host range mutants and semitemperant mutants of T3. *Virology* **3**:527-553.
126. Fröhlich, B., A. Powling, and R. Knippers. 1975. Formation of concatemeric DNA in bacteriophage T7-infected bacteria. *Virology* **65**:455-468.
127. Fuchs, E. 1977. The formation of bacteriophage T7 and T3 lysozymes from inactive precursors. *FEBS Lett.* **80**:27-29.
128. Fujisawa, H., J. I. Miyazaki, and T. Minagawa. 1978. In vitro packaging of phage T3 DNA. *Virology* **87**:394-400.
129. Fujisawa, H., M. Yamagishi, and T. Minagawa. 1980. In vitro formation of the concatemeric DNA of bacteriophage T3 and its biological activity in the in vitro packaging reaction. *Virology* **101**:327-334.
130. Gallant, J. A. 1979. Stringent control in *E. coli*. *Annu. Rev. Genet.* **13**:393-415.
131. Geffer, M., R. Hausmann, M. Gold, and J. Hurwitz. 1966. The enzymatic methylation of RNA and DNA. X. Bacteriophage T3-induced S-adenosylmethionine cleavage. *J. Biol. Chem.* **241**:1995-2005.
132. Geffer, M. L. 1975. DNA replication. *Annu. Rev. Biochem.* **44**:45-78.
133. Geier, G. E., and P. Modrich. 1979. Recognition sequence of the *dam* methylase of *E. coli* K12 and mode of cleavage of *DpnI* endonuclease. *J. Biol. Chem.* **254**:1408-1413.
134. Geier, M. R., and C. R. Merrill. 1972. Lambda phage transcription in human fibroblasts. *Virology* **47**:638-643.
135. Gelinas, R. E., P. A. Myers, and R. J. Roberts. 1977. Two sequence-specific endonucleases from *Moraxella bovis*. *J. Mol. Biol.* **114**:169-179.
136. Gilbert, W. 1976. Starting and stopping sequences for the RNA polymerase, p. 193-205. In R. Losick and M. Chamberlin (ed.), RNA polymerase. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
137. Glickman, B., P. van den Elsten, and M. Radman. 1978. Induced mutagenesis in *dam*<sup>-</sup> mutants of *E. coli*: a role for 6-methyladenine residues in mutation avoidance. *Mol. Gen. Genet.* **183**:307-312.
138. Glover, S. W., J. Schell, N. Symonds, and K. A. Stacey. 1963. The control of host-induced modification by phage P1. *Genet. Res.* **4**:480-482.
139. Goebel, W., and W. Schiess. 1975. The fate of a bacterial plasmid in mammalian cells. *Mol. Gen. Genet.* **138**:213-223.
140. Goebel, W. F., and M. A. Jesaitis. 1952. The somatic antigen of a phage-resistant variant of phase II *Shigella sonnei*. *J. Exp. Med.* **96**:425-438.
141. Gold, M., M. Geffer, R. Hausmann, and J. Hurwitz. 1966. Methylation of DNA. *J. Gen. Physiol.* **49**(no. 6, Part 2):5-28.
142. Gold, M., R. Hausmann, U. Maitra, and J. Hurwitz. 1964. The enzymatic methylation of RNA and DNA. VIII. Effects of bacteriophage infection on the activity of the methylating enzymes. *Proc. Natl. Acad. Sci. U.S.A.* **52**:292-297.
143. Golomb, M., and M. Chamberlin. 1974. Characterization of T7-specific RNA polymerase. III. Resolution of the major in vitro transcripts by gel electrophoresis. *J. Biol. Chem.* **249**:2858-2863.
144. Golomb, M., and H. Chamberlin. 1974. A preliminary map of the major transcription units read by T7 RNA polymerase on the T7 and T3 bacteriophage chromosomes. *Proc. Natl. Acad. Sci. U.S.A.* **71**:760-764.
145. Golomb, M., and M. J. Chamberlin. 1977. T7- and T3-specific RNA polymerases: characterization and mapping of the in vitro transcripts read from T3 DNA. *J. Virol.* **21**:743-752.
146. Gomez, B., and L. Nualart. 1977. Requirement of the bacteriophage T7 0.7 gene for phage growth in the presence of the ColIb factor. *J. Gen. Virol.* **35**:99-106.
147. Gordon, R. L., P. Humphries, and D. J. McConnell. 1978. Restriction enzyme cleavage mapping of T7 virus early region. *Mol. Gen. Genet.* **162**:329-339.
148. Grachev, M. A., E. F. Zaichikov, V. V. Kravchenko, and A. G. Pletnev. 1978. Isolation of promoter and terminator fragments from restriction endonuclease *BsuR*-digested phage T7 DNA. *Dokl. Akad. Nauk. SSSR* **239**:475-478.
149. Grasso, R. J., and K. Paigen. 1969. Loss of host-controlled restriction and modification of phage lambda in *E. coli* K12 previously infected with UV-irradiated coliphage T3. *Virology* **38**:193-194.
150. Grippo, P., and C. C. Richardson. 1971. DNA polymerase of bacteriophage T7. *J. Biol. Chem.* **246**:6867-6873.

151. **Haberman, A.** 1974. The bacteriophage P1 restriction endonuclease. *J. Mol. Biol.* **89**:545-563.
152. **Hadi, S. M., B. Bächli, J. C. W. Shepherd, R. Yuan, K. Ineichen, and T. A. Bickle.** 1979. DNA recognition and cleavage by the EcoP15 restriction endonuclease. *J. Mol. Biol.* **134**: 655-666.
153. **Hadi, S. M., T. A. Bickle, and R. Yuan.** 1975. The role of S-adenosylmethionine in the cleavage of DNA by the restriction endonuclease from *E. coli* K. *J. Biol. Chem.* **250**:4159-4164.
154. **Hagen, F., and E. T. Young.** 1973. Regulation of synthesis of bacteriophage T7 lysozyme mRNA. *Virology* **55**:231-241.
155. **Hagen, F. S., and E. T. Young.** 1974. Preparative polyacrylamide gel electrophoresis of RNA. Identification of multiple molecular species of bacteriophage T7 lysozyme messenger RNA. *Biochemistry* **13**:3394-3400.
156. **Hagen, F. S., and E. T. Young.** 1978. Effect of RNase III on the size of bacteriophage T7 lysozyme mRNA. *J. Virol.* **26**:783-792.
157. **Hagen, F. S., and E. T. Young.** 1978. Effect of RNase III on efficiency of translation of bacteriophage T7 lysozyme mRNA. *J. Virol.* **26**: 793-804.
158. **Hancock, R. E. W., and P. Reeves.** 1975. Bacteriophage resistance in *Escherichia coli* K-12: general pattern of resistance. *J. Bacteriol.* **121**: 983-993.
159. **Hancock, R. E. W., and P. Reeves.** 1976. Lipopolysaccharide-deficient, bacteriophage-resistant mutants of *Escherichia coli* K-12. *J. Bacteriol.* **127**:98-108.
160. **Hattman, S., J. E. Brooks, and M. Masupekar.** 1978. Sequence specificity of the P1 modification methylase (M.EcoP1) and the DNA methylase (M.Ecodam) controlled by the *E. coli* *dam* gene. *J. Mol. Biol.* **126**:367-380.
161. **Hausmann, R.** 1967. Synthesis of an S-adenosylmethionine-cleaving enzyme in T3-infected *Escherichia coli* and its disturbance by coinfection with enzymatically incompetent bacteriophage. *J. Virol.* **1**:57-63.
162. **Hausmann, R.** 1968. Sedimentation analysis of phage T7-directed DNA synthesized in the presence of a dominant conditional lethal phage gene. *Biochem. Biophys. Res. Commun.* **31**:609-615.
163. **Hausmann, R.** 1973. Host-dependent gene expression of bacteriophage T3 and T7, p. 263-276. In B. A. Hamkalo and J. Papaconstantinou (ed.), *Molecular cytogenetics*. Plenum Publishing Corp., New York.
164. **Hausmann, R.** 1973. The genetics of T-odd phages. *Annu. Rev. Microbiol.* **27**:51-68.
165. **Hausmann, R.** 1976. Bacteriophage T7 genetics. *Curr. Top. Microbiol. Immunol.* **75**:77-110.
166. **Hausmann, R., and B. Gomez.** 1967. Amber mutants of bacteriophages T3 and T7 defective in phage-directed DNA synthesis. *J. Virol.* **1**: 779-792.
167. **Hausmann, R., and B. Gomez.** 1968. Bacteriophage T3- and T7-directed deoxyribonucleases. *J. Virol.* **2**:265-266.
168. **Hausmann, R., and E. Härle.** 1971. Expression of the genomes of the related phages T3 and T7, p. 467-498. In E. Broda et al. (ed.), *Proceedings of the First European Biophysics Congress*, Baden, vol. 1. Wiener Med. Akademie, Vienna.
169. **Hausmann, R., and K. LaRue.** 1969. Variations in sedimentation patterns among deoxyribonucleic acids synthesized after infection of *Escherichia coli* by different amber mutants of bacteriophage T7. *J. Virol.* **3**:278-281.
170. **Hausmann, R., and C. Tomkiewicz.** 1976. Genetic analysis of template specificity of RNA polymerases (gene 1 products) coded by phage T3 x T7 recombinants within gene 1, p. 731-743. In R. Losick and M. Chamberlin (eds.), *RNA polymerase*. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
171. **Hausmann, R. L., E. P. Almeida-Magalhaes, and C. Aranjó.** 1961. Isolation and characterization of hybrids between bacteriophage T3 and T7. *An. Microbiol.* **9**:511-526.
172. **Hausmann, R. L., E. P. Almeida Magalhaes, and C. Aranjó.** 1962. Serological properties of two related coliphages, T7 and Cro. *An. Microbiol.* **10**:35-42.
173. **Hausmann, R. L., E. P. Almeida-Magalhaes, and C. Aranjó.** 1962. Intracellular compatibility of coliphages T3, T7 and Cro. *An. Microbiol.* **10**:43-62.
174. **Hayes, W.** 1968. The genetics of bacteria and their viruses, 2nd ed. Blackwell Scientific Publications, Ltd., Oxford.
175. **Helland, D., and A. P. Nygaard.** 1975. Selected genes of T7 DNA associated with the *E. coli* membrane following phage infection. *FEBS Lett.* **50**:13-16.
176. **Henner, W. D., I. Kleber, and K. Benzinger.** 1973. Transfection of *Escherichia coli* spheroplasts. II. Facilitation of transfection and stabilization of spheroplasts by different basic polymers. *J. Virol.* **12**:741-747.
177. **Hercules, K., S. Jovanovich, and W. Sauerbier.** 1976. Early gene expression in bacteriophage T7. I. In vivo synthesis, inactivation, and translational utilization of early mRNA's. *J. Virol.* **17**:642-658.
178. **Hercules, K., M. Schweiger, and W. Sauerbier.** 1974. Cleavage by RNase III converts T3 and T7 early precursor RNA into translatable message. *Proc. Natl. Acad. Sci. U.S.A.* **71**:840-844.
179. **Herrlich, P., H. J. Rahmsdorf, S. H. Pai, and M. Schweiger.** 1974. Translational control induced by bacteriophage T7. *Proc. Natl. Acad. Sci. U.S.A.* **71**:1088-1092.
180. **Herrlich, P., M. Schweiger, and W. Sauerbier.** 1971. Host- and phage-RNA polymerase mediated synthesis of T7 lysozyme in vivo. *Mol. Gen. Genet.* **112**:152-160.
181. **Hershey, A. D. (ed.).** 1971. The bacteriophage lambda. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
182. **Hershey, A. D., A. Garen, D. K. Fraser, and J. D. Hudis.** 1954. Growth and inheritance in

- bacteriophage. Carnegie Inst. Washington Yearb. 53:210-225.
183. Hesselbach, B. A., and D. Nakada. 1975. Inactive complex formation between *E. coli* RNA polymerase and inhibitor protein purified from T7 phage infected cells. *Nature (London)* 258:354-357.
184. Hesselbach, B. A., and D. Nakada. 1977. "Host shutoff" function of bacteriophage T7: involvement of T7 gene 2 and gene 0.7 in the inactivation of *E. coli* RNA polymerase. *J. Virol.* 24:736-745.
185. Hesselbach, B. A., and D. Nakada. 1977. I protein: bacteriophage T7-coded inhibitor of *Escherichia coli* RNA polymerase. *J. Virol.* 24:746-760.
186. Hesselbach, B. A., Y. Yamada, and D. Nakada. 1974. Isolation of an inhibitor protein of *E. coli* RNA polymerase from T7 phage infected cell. *Nature (London)* 252:71-74.
187. Hiebsch, R., and M. S. Center. 1977. Intracellular organization of bacteriophage T7 DNA: analysis of parental bacteriophage T7 DNA-membrane and DNA-protein complexes. *J. Virol.* 22:540-547.
188. Hillenbrand, G., G. Morelli, E. Lanka, and E. Scherzinger. 1979. Bacteriophage T7 DNA primase: a multifunctional enzyme involved in DNA replication. *Cold Spring Harbor Symp. Quant. Biol.* 43:449-459.
189. Hinkle, D. C. 1980. Evidence for direct involvement of T7 RNA polymerase in bacteriophage DNA replication. *J. Virol.* 34:136-141.
190. Hinkle, D. C., and M. J. Chamberlin. 1972. Studies on the binding of *E. coli* RNA polymerase to DNA. I. The role of sigma subunit in site selection. *J. Mol. Biol.* 70:157-185.
191. Hinkle, D. C., and C. C. Richardson. 1974. Bacteriophage T7 DNA replication in vitro. Requirements for DNA synthesis and characterization of the product. *J. Biol. Chem.* 249:2974-2984.
192. Hinkle, D. C., and C. C. Richardson. 1975. Bacteriophage T7 DNA replication in vitro. Purification and properties of the gene 4 product of bacteriophage T7. *J. Biol. Chem.* 250:5523-5529.
193. Hirsch-Kaufmann, M., M. Pfennig-Yeh, H. Ponta, P. Herrlich, and M. Schweiger. 1976. A virus-specified mechanism for the prevention of multiple infection: T7 and T3 mutual and superinfection exclusion. *Mol. Genet.* 149:243-249.
194. Holliday, R., and J. E. Pugh. 1975. DNA modification mechanisms and gene activity during development. *Science* 187:226-232.
195. Holmgren, A., I. Ohlsson, and M. L. Grankvist. 1978. Thioredoxin from *E. coli*. Radioimmunological and enzymatic determinations in wild type cells and mutants defective in phage T7 DNA replication. *J. Biol. Chem.* 253:430-436.
196. Hopper, J. E., G. Ko, and E. T. Young. 1975. Comparative analysis of the in vivo and in vitro expression of bacteriophage T7 messenger RNAs during infection of *E. coli*. *J. Mol. Biol.* 94:539-554.
197. Hori, K., D. F. Mark, and C. C. Richardson. 1979. DNA polymerase of bacteriophage T7. Purification and properties of the phage encoded subunit, the gene 5 protein. *J. Biol. Chem.* 254:11591-11597.
198. Hori, K., D. F. Mark, and C. C. Richardson. 1979. DNA polymerase of bacteriophage T7. Characterization of the exonuclease activities of the gene 5 protein and the reconstituted polymerase. *J. Biol. Chem.* 254:11598-11604.
199. Hotchkiss, R. D. 1974. Models of genetic recombination. *Annu. Rev. Microbiol.* 28:445-468.
200. Hsieh, T., and J. C. Wang. 1976. Physicochemical studies on interactions between DNA and RNA polymerase. Isolation and mapping of a T7 fragment containing the early promoters for *E. coli* RNA polymerase. *Biochemistry* 15:5776-5783.
201. Humphries, P., R. L. Gordon, D. J. McConnell, and P. Connolly. 1974. Endonuclease *R.Hind* fragments of T7 DNA. *Virology* 58:25-31.
202. Humphries, P., D. J. McConnell, and P. Connolly. 1974. Interaction of *E. coli* DNA-dependent RNA polymerase and endonuclease *R.Hind* fragments of T7 DNA. *Biochim. Biophys. Acta* 366:70-78.
203. Hyman, R. W., I. Brunovskis, and W. C. Summers. 1974. A biochemical comparison of the related bacteriophage T7,  $\phi$ I,  $\phi$ II, W31, H, and T3. *Virology* 57:189-206.
204. Ihler, G. M., and C. A. Thomas. 1970. Equal incorporation of both parental T7 DNA strands into intracellular concatemeric DNA. *J. Virol.* 6:877-880.
205. Inouye, M., N. Arhneim, and R. Sternglanz. 1973. Bacteriophage T7 lysozyme is an N-acetylmuramyl-L-alanine amidase. *J. Biol. Chem.* 248:7247-7252.
206. Issinger, O. G., H. Beier, and R. Hausmann. 1973. In vivo and in vitro "phenotypic mixing" with amber mutants of phages T3 and T7. *Mol. Gen. Genet.* 122:81-88.
207. Issinger, O. G., and H. Falk. 1976. Comparative studies on the structural proteins of T3 and T7 phages. *Arch. Virol.* 52:217-231.
208. Itoh, T., and J. I. Tomizawa. 1977. Involvement of DNA gyrase in bacteriophage T7 DNA replication. *Nature (London)* 270:78-80.
209. Iyer, V. N., and A. P. James. 1978. Single-cell studies on the carrier state of bacteriophage 1Ke, a virus specific for conjugative plasmids of the N incompatibility and conjugative group. *Can. J. Microbiol.* 24:1595-1601.
210. Jensen, H. B., and I. F. Pryme. 1974. The separation of a gene 3.5 directed activity and a lytic activity ("lysozyme") in T7-infected *E. coli* B cells. *Biochem. Biophys. Res. Commun.* 59:1117-1123.
211. Jovin, T. M. 1976. Recognition mechanisms of DNA-specific enzymes. *Annu. Rev. Biochem.* 45:889-920.
212. Kadesch, T. R., R. C. Williams, and M. J.



- Chamberlin. 1980. Electron microscopic studies of the binding of *E. coli* RNA polymerase to DNA. I. Characterization of the non-specific interactions of holoenzyme with a restriction fragment of bacteriophage T7 DNA. *J. Mol. Biol.* **136**:65-78.
213. Kadesch, T. R., R. C. Williams, and M. J. Chamberlin. 1980. Electron microscopic studies of the binding of *E. coli* RNA polymerase to DNA. II. Formation of multiple promoter-like complexes at non-promoter sites. *J. Mol. Biol.* **136**:79-93.
214. Kan, N. C., J. A. Lautenberger, M. H. Edgell, and C. A. Hutchinson. 1979. The nucleotide sequence recognized by the *E. coli* K12 restriction and modification enzymes. *J. Mol. Biol.* **130**:191-209.
215. Kao, P. C., J. D. Regan, and E. Volkin. 1973. Fate of homologous and heterologous DNAs after incorporation into human skin fibroblasts. *Biochim. Biophys. Acta* **324**:1-13.
216. Karu, A. E., Y. Sakaki, H. Echols, and S. Linn. 1975. The gamma protein specified by bacteriophage lambda. Structure and inhibitory activity for the *recBC* enzyme of *E. coli*. *J. Biol. Chem.* **250**:7377-7387.
217. Kassavetis, G. A., and M. J. Chamberlin. 1979. Mapping of class II promoter sites utilized in vitro by T7-specific RNA polymerases on bacteriophage T7 DNA. *J. Virol.* **29**:196-208.
218. Kassavetis, G. A., K. M. Kaya, and M. J. Chamberlin. 1978. *E. coli* RNA polymerase-rifampicin complexes bound at promoter sites block RNA chain elongation by *E. coli* RNA polymerase and T7-specific RNA polymerase. *Biochemistry* **17**:5798-5804.
219. Kawakami, K., T. Saitoh, and A. Ishihama. 1979. Biosynthesis of RNA polymerase in *E. coli*. IX. Growth-dependent variations in the synthesis rate, content, and distribution of RNA polymerase. *Mol. Gen. Genet.* **174**:107-116.
220. Kelly, T. J., and H. D. Smith. 1970. A restriction enzyme from *Hemophilus influenzae*. II. Base sequence of recognition site. *J. Mol. Biol.* **51**:393-409.
221. Kelly, T. J., and C. A. Thomas. 1969. An intermediate in the replication of bacteriophage T7 DNA molecules. *J. Mol. Biol.* **44**:459-475.
222. Kerr, C., and P. D. Sadowski. 1972. Gene 6 exonuclease of bacteriophage T7. I. Purification and properties of the enzyme. *J. Biol. Chem.* **247**:305-310.
223. Kerr, C., and P. D. Sadowski. 1972. Gene 6 exonuclease of bacteriophage T7. II. Mechanism of the reaction. *J. Biol. Chem.* **247**:311-318.
224. Kerr, C., and P. D. Sadowski. 1974. Packaging and maturation of DNA of bacteriophage T7 in vitro. *Proc. Natl. Acad. Sci. U.S.A.* **71**:3545-3549.
225. Kerr, C., and P. Sadowski. 1975. The involvement of genes 3, 4, 5 and 6 in genetic recombination of bacteriophage T7. *Virology* **65**:281-285.
226. Kiefer, M., N. Neff, and M. J. Chamberlin. 1977. Transcriptional termination at the end of the early region of bacteriophages T3 and T7 is not affected by polarity suppressors. *J. Virol.* **22**:548-552.
227. Klein, A. 1965. Mechanismen der wirtskontrollierten Modifikation des Phagen TI. *Z. Vererbungsl.* **96**:346-363.
228. Kleppe, G., H. B. Jensen, and I. F. Pryme. 1977. Purification and characterization of the lytic enzyme N-acetylmuramyl-L-alanine amidase of bacteriophage T7. *Eur. J. Biochem.* **76**:317-326.
229. Knippers, R. 1979. T7 DNA replication—a brief introduction. *FEBS (Fed. Eur. Biochem. Soc.) Proc. Meet.* **51**:331-335.
230. Knippers, R., W. Strätling, and E. Krause. 1973. Some biochemical elements in bacteriophage T7 DNA replication in vitro, p. 451-461. In R. D. Wells and R. B. Inman (ed.), *DNA synthesis in vitro*. Medical and Technical Publishing Co., Ltd., Lancaster, England.
231. Koller, T., O. Kübler, R. Portmann, and J. M. Sogo. 1978. High resolution physical mapping of specific binding sites of *E. coli* RNA polymerase on the DNA of bacteriophage T7. *J. Mol. Biol.* **120**:121-131.
232. Koller, T., J. M. Sogo, and H. Bujard. 1974. An electron microscopic method for studying nucleic acid-protein complexes. Visualization of RNA polymerase bound to the DNA of bacteriophages T3 and T7. *Biopolymers* **13**:995-1009.
233. Kolodner, R., Y. Masamune, J. E. LeClerc, and C. C. Richardson. 1978. Gene 4 protein of bacteriophage T7. Purification, physical properties, and stimulation of T7 DNA polymerase during the elongation of polynucleotide chains. *J. Biol. Chem.* **253**:566-573.
234. Kolodner, R., and C. C. Richardson. 1977. Replication of duplex DNA by bacteriophage T7 DNA polymerase and gene 4 protein is accompanied by hydrolysis of nucleoside 5'-triphosphates. *Proc. Natl. Acad. Sci. U.S.A.* **74**:1525-1529.
235. Kolodner, R., and C. C. Richardson. 1978. Gene 4 protein of bacteriophage T7. Characterization of the product synthesized by the T7 DNA polymerase and gene 4 protein in the absence of ribonucleoside 5'-triphosphates. *J. Biol. Chem.* **253**:574-584.
236. Kolter, R., and D. Helinski. 1979. Regulation of initiation of DNA replication. *Annu. Rev. Genet.* **13**:355-391.
237. Kornberg, A. 1979. Aspects of DNA replication. *Cold Spring Harbor Symp. Quant. Biol.* **43**:1-9.
238. Korobko, V. G., S. A. Tchuvpilo, and M. N. Kolosov. 1980. Nucleotide sequence of phage T7 gene 0.4. *Bioorg. Khim.* **6**:1114-1116. (In Russian.)
239. Korsten, K. H., C. Tomkiewicz, and R. Hausmann. 1979. The strategy of infection as a

- criterion for phylogenetic relationship of non-coli phages morphologically similar to phage T7. *J. Gen. Virol.* **43**:57-73.
240. **Kramer, R. A., and M. Rosenberg.** 1976. The isolation and characterization of bacteriophage T7 mRNA fragments containing an RNaseIII cleavage site. *Nucleic Acids Res.* **3**:2411-2426.
241. **Kramer, R. A., M. Rosenberg, and J. A. Steitz.** 1974. Nucleotide sequences of the 5' and 3' termini of bacteriophage T7 early mRNA synthesized in vivo: evidence for sequence specificity of mRNA processing. *J. Mol. Biol.* **89**:767-776.
242. **Kreuzer, K. N., and N. R. Cozzarelli.** 1979. *Escherichia coli* mutants thermosensitive for deoxyribonucleic acid gyrase subunit A: effects on deoxyribonucleic acid replication, transcription, and bacteriophage growth. *J. Bacteriol.* **140**:424-435.
243. **Krueger, D. H., W. Presber, S. Hansen, and H. A. Rosenthal.** 1975. Biological functions of the bacteriophage T3 SAMase gene. *J. Virol.* **16**:453-455.
244. **Krüger, D. H., L. S. Chernin, S. Hansen, H. A. Rosenthal, and D. M. Goldfarb.** 1978. Protection of foreign DNA against host-controlled restriction in bacterial cells. I. Protection of F' plasmid DNA by preinfecting with bacteriophages T3 or T7. *Mol. Gen. Genet.* **159**:107-110.
245. **Krüger, D. H., G. Gola, I. Weissshuhn, and S. Hansen.** 1978. The *ocr* gene function of bacterial viruses T3 and T7 prevents host-controlled modification. *J. Gen. Virol.* **41**:189-192.
246. **Krüger, D. H., S. Hansen, W. Presber, M. Rudolph, D. Scholz, and H. A. Rosenthal.** 1979. Wirtsbereichsverhalten der Bakteriophagen T3 und T7 an *E. coli* K12-Stämmen. *Z. Allg. Mikrobiol.* **19**:473-480.
247. **Krüger, D. H., S. Hansen, W. Presber, and C. Schroeder.** 1978. Host-dependent modification of virus strains affecting their adsorption ability, p. 543-545. In L. Shindarov (ed.), *Proceedings of the Seventh International Congress of Infectious and Parasitic Diseases*, vol. 3. Union of Scientific Medical Societies in Bulgaria, Varna.
248. **Krüger, D. H., S. Hansen, and C. Schroeder.** 1981. Host-dependent modification of bacterial virus T3 affecting its adsorption ability. *Virology* **102**:444-446.
249. **Krüger, D. H., S. Hansen, and C. Schroeder.** 1981. Virus adaptation to host cells: the non-classical modification of phage T3. *Z. Allg. Mikrobiol.* **20**:495-502.
250. **Krüger, D. H., S. Hansen, C. Schroeder, and W. Presber.** 1977. Host-dependent modification of bacteriophage T7 and SAMase-negative T3 derivatives affecting their adsorption ability. *Mol. Gen. Genet.* **153**:107-110.
251. **Krüger, D. H., W. Presber, S. Hansen, and H. A. Rosenthal.** 1977. Different restriction of bacteriophages T3 and T7 by PI-lysogenic cells and the role of the T3-coded SAMase. *Z. Allg. Mikrobiol.* **17**:581-591.
252. **Krüger, D. H., W. Presber, W. Mann, and H. A. Rosenthal.** 1972. Untersuchungen an T3-Phagen. III. Über einen unterschiedlichen Nährstoffmangel-effekt auf die Vermehrung der Phagen T3 und T7. *Acta Biol. Med. Germ.* **29**:477-482.
253. **Krüger, D. H., D. Scholz, and H. A. Rosenthal.** 1975. Untersuchungen an T3-Phagen. VII. Cross streak-Komplementierung von temperatursensitiven Mutanten des Bakteriophagen T3 unter konstant nonpermissiven Bedingungen. *Z. Allg. Mikrobiol.* **15**:81-88.
254. **Krüger, D. H., C. Schroeder, S. Hansen, and H. A. Rosenthal.** 1977. Active protection by bacteriophages T3 and T7 against *E. coli* Band K-specific restriction of their DNA. *Mol. Gen. Genet.* **153**:99-106.
255. **Kuemmerle, N. B., and W. E. Masker.** 1977. In vitro packaging of UV radiation-damaged DNA from bacteriophage T7. *J. Virol.* **23**:509-516.
256. **Kuhn, B., M. Abdel-Monem, and H. Hoffmann-Berling.** 1979. DNA helicases. *Cold Spring Harbor Symp. Quant. Biol.* **43**:63-67.
257. **Lacks, S., and B. Greenberg.** 1975. A deoxyribonuclease of *Diplococcus pneumoniae* specific for methylated DNA. *J. Biol. Chem.* **250**:4060-4066.
258. **Lacks, S., and B. Greenberg.** 1977. Complementary specificity of restriction endonucleases of *Diplococcus pneumoniae* with respect to DNA methylation. *J. Mol. Biol.* **114**:153-168.
259. **Landy, A., E. Ruedisueli, L. Robinson, C. Foeller, and W. Ross.** 1974. Digestion of DNAs from bacteriophage T7, lambda, and  $\phi$ 80h with site-specific nucleases from *Hemophilus influenzae* strain Rc and strain Rd. *Biochemistry* **13**:2134-2142.
260. **Langman, L., and V. Paetkau.** 1978. Purification and structures of recombining and replicating bacteriophage T7 DNA. *J. Virol.* **25**:562-569.
261. **Lark, C.** 1979. Methylation-dependent DNA synthesis in *Escherichia coli* mediated by DNA polymerase I. *J. Bacteriol.* **137**:44-50.
262. **Lathe, R.** 1978. RNA polymerase of *E. coli*. *Curr. Top. Microbiol. Immunol.* **83**:37-91.
263. **Laurent, T. C., S. C. Moore, and P. Reichard.** 1964. Enzymatic synthesis of deoxyribonucleotides. IV. Isolation and characterization of thioredoxin, the hydrogen donor from *E. coli* B. *J. Biol. Chem.* **239**:3436-3444.
264. **Lautenberger, J. A., N. C. Kan, D. Lackey, S. Linn, M. H. Edgell, and C. A. Hutchinson.** 1978. Recognition site of *E. coli* B restriction enzyme on  $\phi$ XsBI and Simian virus 40 DNAs: an interrupted sequence. *Proc. Natl. Acad. Sci. U.S.A.* **75**:2271-2275.
265. **Lawhorne, L., I. Kleber, C. Mitchell, and R. Benzinger.** 1973. Transfection of *Escherichia coli* spheroplasts. II. Relative infectivity of native, denatured, and renatured lambda, T7, T5, T4, and P22 bacteriophage DNAs. *J. Virol.* **12**:733-740.
266. **Leavitt, J. C., L. M. Schlechtman, P. O. P.**

- Tso, E. Rorenfreund, and A. Bendich. 1976. The attachment and penetration of T7 DNA/phage in Syrian hamster embryonic cells. *Biochim. Biophys. Acta* **435**:167-183.
267. LeClerc, J. E., and C. C. Richardson. 1979. Bacteriophage T7 DNA replication in vitro. XVI. Gene 2 protein of bacteriophage T7: purification and requirement for packaging of T7 DNA in vitro. *Proc. Natl. Acad. Sci. U.S.A.* **76**: 4852-4856.
268. Lederberg, S. 1957. Suppression of the multiplication of heterologous bacteriophages in lysogenic bacteria. *Virology* **3**:496-513.
269. Lee, M., and R. C. Miller. 1974. T7 exonuclease (gene 6) is necessary for molecular recombination of bacteriophage T7. *J. Virol.* **14**:1040-1048.
270. Li, L. I., V. I. Tanyashin, N. I. Matvienko, and A. A. Bayev. 1975. Production of specific fragments of T4 DNA by endonuclease *EcoRI*. *Dokl. Akad. Nauk SSSR* **223**:1262-1265. (In Russian.)
271. Lindberg, A. A. 1973. Bacteriophage receptors. *Annu. Rev. Microbiol.* **27**:205-241.
272. Linn, S. 1978. The 1978 Nobel prize in physiology and medicine. *Science* **202**:1069-1071.
273. Lippincott, J. A. 1977. Molecular basis of plant tumor induction. *Nature (London)* **269**:465-466.
274. Losick, R., and J. Pero. 1976. *Bacillus subtilis* RNA polymerase and its modification in sporulating and phage-infected bacteria. *Adv. Enzymol. Relat. Areas Mol. Biol.* **44**:165-185.
275. Ludwig, R. A., and W. C. Summers. 1975. A restriction fragment analysis of the T7 left-early region. *Virology* **68**:360-373.
276. Ludwig, R. A., and W. C. Summers. 1976. Localization of RNA polymerase binding sites on T7 DNA. *Virology* **71**:278-290.
277. Luria, S. E. 1953. Host-induced modification of viruses. *Cold Spring Harbor Symp. Quant. Biol.* **18**:237-244.
278. Lwoff, A. 1953. Lysogeny. *Bacteriol. Rev.* **17**: 269-337.
279. Mahadik, S. P., B. Dharmgrongartama, and P. R. Srinivasan. 1972. An inhibitory protein of *E. coli* RNA polymerase in bacteriophage T3-infected cells. *Proc. Natl. Acad. Sci. U.S.A.* **69**:162-166.
280. Mahadik, S. P., B. Dharmgrongartama, and P. R. Srinivasan. 1974. Regulation of host RNA synthesis in bacteriophage T3-infected cells. *J. Biol. Chem.* **249**:1787-1791.
281. Maitra, U. 1971. Induction of a new RNA polymerase in *E. coli* infected with bacteriophage T3. *Biochem. Biophys. Res. Commun.* **43**:443-450.
282. Maitra, U., and H. H. Huang. 1972. Initiation, release, and reinitiation of RNA chains by bacteriophage T3-induced polymerase from T3 templates. *Proc. Natl. Acad. Sci. U.S.A.* **69**:55-59.
283. Maitra, U., R. A. Salvo, and P. R. Chakraborty. 1974. Specificity of RNA chain initiation by bacteriophage T3-induced RNA polymerase. *J. Biol. Chem.* **249**:5835-5839.
284. Majumder, H. K., S. Bishayee, P. R. Chakraborty, and U. Maitra. 1977. RNaseIII cleavage of bacteriophage T3 RNA polymerase transcripts to late T3 mRNAs. *Proc. Natl. Acad. Sci. U.S.A.* **74**:4891-4894.
285. Majumder, H. K., U. Maitra, and M. Rosenberg. 1979. Termination of transcription by bacteriophage T3 RNA polymerase: homogeneous 3'-terminal oligonucleotide sequence of in vitro RNA polymerase transcripts. *Proc. Natl. Acad. Sci. U.S.A.* **76**:5110-5113.
286. Mäkelä, O., P. H. Mäkelä, and S. Soikkeli. 1964. Sex-specificity of bacteriophage T7. *Ann. Med. Exp. Biol. Fenn.* **42**:188-195.
287. Makino, O., F. Kawamura, H. Saito, and Y. Ikeda. 1979. Inactivation of restriction endonuclease *BamNI* after infection with phage  $\phi$ NR2. *Nature (London)* **277**:64-66.
288. Male, C. J., and J. R. Christensen. 1970. Synthesis of messenger RNA after bacteriophage T1 infection. *J. Virol.* **6**:727-737.
289. Mangel, W. F., and M. J. Chamberlin. 1974. Studies of RNA chain initiation by *E. coli* RNA polymerase bound to T7 DNA. I. An assay for the rate and extent of RNA chain initiation. *J. Biol. Chem.* **249**:2995-3001.
290. Mann, W., S. Michel, B. Brux, S. Elgammal, and G. Richter. 1972. Replication of T3-phages in a temperature-sensitive mutant of *E. coli* (*E. coli* CRT266). *Stud. Biophys.* **31/32**: 15-21.
291. Marinus, M. G., and N. R. Morris. 1973. Isolation of deoxyribonucleic acid methylase mutants of *Escherichia coli* K-12. *J. Bacteriol.* **114**:1143-1150.
292. Mark, D. F., and C. C. Richardson. 1976. *E. coli* thioredoxin: a subunit of bacteriophage T7 DNA polymerase. *Proc. Natl. Acad. Sci. U.S.A.* **73**:780-784.
293. Masamuno, Y., G. D. Frenkel, and C. C. Richardson. 1971. A mutant of bacteriophage T7 deficient in polynucleotide ligase. *J. Biol. Chem.* **246**:6847-6879.
294. Masker, W. E., and N. B. Kuemmerle. 1980. In vitro recombination of bacteriophage T7 DNA damaged by UV radiation. *J. Virol.* **33**: 330-339.
295. Masker, W. E., N. B. Kuemmerle, and D. P. Allison. 1978. In vitro packaging of bacteriophage T7 DNA synthesized in vitro. *J. Virol.* **27**:149-163.
296. Masker, W. E., and C. C. Richardson. 1976. Bacteriophage T7 DNA replication in vitro. V. Synthesis of intact chromosomes of bacteriophage T7. *J. Mol. Biol.* **100**:543-556.
297. Masker, W. E., and C. C. Richardson. 1976. Bacteriophage T7 DNA replication in vitro. VI. Synthesis of biologically active T7 DNA. *J. Mol. Biol.* **100**:557-567.
298. McAllister, W. T., and C. L. Barrett. 1977. Roles of the early genes of bacteriophage T7 in shutoff of host macromolecular synthesis. *J. Virol.* **23**:543-553.
299. McAllister, W. T., and C. L. Barrett. 1977. Superinfection exclusion by bacteriophage T7.

- J. Virol. **24**:709-711.
300. McAllister, W. T., and C. L. Barrett. 1977. Hybridization mapping of restriction fragments from the early region of bacteriophage T7 DNA. *Virology* **82**:275-287.
301. McAllister, W. T., H. Küpper, and E. K. F. Bautz. 1973. Kinetics of the transcription by the bacteriophage T3 RNA polymerase in vitro. *Eur. J. Biochem.* **34**:489-501.
302. McAllister, W. T., and R. J. McCarron. 1977. Hybridization of the in vitro products of bacteriophage T7 RNA polymerase to restriction fragments of T7 DNA. *Virology* **82**:288-298.
303. McAllister, W. T., and H. L. Wu. 1978. Regulation of transcription of the late genes of bacteriophage T7. *Proc. Natl. Acad. Sci. U.S.A.* **75**:804-808.
304. McConnell, D. J. 1979. The DNA sequence at the T7 C promoter. *Nucleic Acids Res.* **6**:525-544.
305. McConnell, D. J. 1979. Control sites in the sequence at the beginning of T7 gene I. *Nucleic Acids Res.* **6**:3491-3503.
306. McCorquodale, D. J. 1975. The T-odd bacteriophages. *Crit. Rev. Microbiol.* **5**:101-159.
307. McDonell, M. W., M. N. Simon, and F. W. Studier. 1977. Analysis of restriction fragments of T7 DNA and determination of molecular weights by electrophoresis in neutral and alkaline gels. *J. Mol. Biol.* **110**:119-146.
308. McGhee, J. D., and G. D. Ginder. 1979. Specific DNA methylation sites in the vicinity of the chicken  $\beta$ -globin genes. *Nature (London)* **280**:419-420.
309. Merrill, C. R. 1974. Bacteriophage interactions with higher organisms. *Trans. N.Y. Acad. Sci.* **36**:265-272.
310. Merrill, C. R. 1977. Interactions of bacterial viruses and bacterial genes with animal systems, p. 83-96. *In* I. Rubinstein et al. (ed.), *Molecular genetic modification of eucaryotes*. Academic Press, Inc., New York.
311. Merrill, C. R., M. R. Geier, and J. C. Petricciani. 1971. Bacterial virus gene expression in human cells. *Nature (London)* **233**:398-400.
312. Milch, H., M. Gyenes, and G. Herman. 1975. Characterization on the basis of phage restriction of R plasmids derived from *E. coli* strains, p. 391-396. *In* S. Mitsuhashi et al. (ed.), *Drug inactivating enzymes and antibiotic resistance*. Avicenum, Nakladatelství, Prague.
313. Miller, R. C. 1972. Asymmetric annealing of an RNA linked DNA molecule isolated during the initiation of bacteriophage T7 DNA replication. *Biochem. Biophys. Res. Commun.* **49**:1082-1086.
314. Miller, R. C. 1975. Replication and molecular recombination of T-phage. *Annu. Rev. Microbiol.* **29**:355-376.
315. Miller, R. C., M. Lee, D. G. Scraba, and V. Paetkau. 1976. The role of bacteriophage T7 exonuclease (gene 6) in genetic recombination and production of concatemers. *J. Mol. Biol.* **101**:223-234.
316. Minkley, E. G. 1974. Transcription of the early region of bacteriophage T7: characterization of the in vivo transcripts. *J. Mol. Biol.* **83**:289-304.
317. Minkley, E. G. 1974. Transcription of the early region of bacteriophage T7: specificity and selectivity in the in vitro initiation of RNA synthesis. *J. Mol. Biol.* **83**:305-331.
318. Minkley, E. G., and D. Pribnow. 1973. Transcription of the early region of bacteriophage T7: selective initiation with dinucleotides. *J. Mol. Biol.* **77**:255-277.
319. Miyazaki, J. I., H. Fujisawa, and T. Minagawa. 1979. Biological activity of purified bacteriophage T3 prohead and prohead-like structures as precursors for in vitro head assembly. *Virology* **91**:283-290.
320. Miyazaki, J. I., Y. Ryo, H. Fujisawa, and T. Minagawa. 1978. Mutation in bacteriophage T3 affecting host cell lysis. *Virology* **89**:327-329.
321. Modrich, P., and C. C. Richardson. 1975. Bacteriophage T7 DNA replication in vitro. A protein of *E. coli* required for bacteriophage T7 DNA polymerase activity. *J. Biol. Chem.* **250**:5508-5514.
322. Modrich, P., and C. C. Richardson. 1975. Bacteriophage T7 DNA replication in vitro. Bacteriophage T7 DNA polymerase: an enzyme composed of phage and host-specified subunits. *J. Biol. Chem.* **250**:5515-5522.
323. Molineux, I. J., S. Friedman, and M. L. Gefter. 1974. Purification and properties of the *E. coli* DNA unwinding protein. Effects on DNA synthesis in vitro. *J. Biol. Chem.* **249**:6090-6098.
324. Molineux, I. J., and M. L. Gefter. 1974. Properties of the *E. coli* DNA binding (unwinding) protein: interaction with DNA polymerase and DNA. *Proc. Natl. Acad. Sci. U.S.A.* **71**:3858-3862.
325. Morrison, T. G., D. D. Blumberg, and M. H. Malamy. 1974. T7 protein synthesis in F' episome-containing cells: assignment of specific proteins to three translational groups. *J. Virol.* **13**:386-393.
326. Murialdo, H., and A. Becker. 1978. Head morphogenesis of complex double-stranded deoxyribonucleic acid bacteriophages. *Microbiol. Rev.* **42**:529-576.
327. Nathans, D. 1979. Restriction endonucleases, simian virus 40, and the new genetics. *Science* **206**:903-909.
328. Nathans, D., and H. O. Smith. 1975. Restriction endonucleases in the analysis and restructuring of DNA molecules. *Annu. Rev. Biochem.* **44**:273-293.
329. Nierlich, D. P. 1978. Regulation of bacterial growth, RNA, and protein synthesis. *Annu. Rev. Microbiol.* **32**:393-432.
330. Nierman, W. C., and M. J. Chamberlin. 1979. Studies of RNA chain initiation by *E. coli* RNA polymerase bound to T7 DNA. Direct analysis of the kinetics and extent of RNA chain initiation at T7 promoter A1. *J. Biol. Chem.* **254**:7921-7926.

331. Nikolskaya, I. I., N. G. Lopatina, N. M. Chaplygina, and S. S. Debov. 1976. The host specificity system in *E. coli* SK. *Mol. Cell. Biochem.* 13:79-87.
332. Niles, E. G., and R. C. Condit. 1975. Translational mapping of bacteriophage T7 RNAs synthesized in vitro by purified T7 RNA polymerase. *J. Mol. Biol.* 98:57-67.
333. Niles, E. G., S. W. Conlon, and W. C. Summers. 1974. Purification and physical characterization of T7 RNA polymerase from T7-infected *E. coli* B. *Biochemistry* 13:3904-3912.
334. Nossal, N. G. 1969. A T4 bacteriophage mutant which lacks DNA polymerase but retains the polymerase-associated nuclease. *J. Biol. Chem.* 244:218-220.
335. Nossal, N. G., and M. S. Herschfield. 1971. Nuclease activity in a fragment of T4 DNA polymerase encoded by the amber mutant amb22. *J. Biol. Chem.* 246:5414-5426.
336. Oakley, J. L., and J. E. Coleman. 1977. Structure of a promoter for T7 RNA polymerase. *Proc. Natl. Acad. Sci. U.S.A.* 74:4266-4270.
337. Oakley, J. L., J. A. Pascale, and J. E. Coleman. 1975. T7 RNA polymerase: conformation, functional groups and promoter binding. *Biochemistry* 14:4684-4691.
338. Oakley, J. L., R. E. Strothcamp, A. H. Sarris, and J. E. Coleman. 1979. T7 RNA polymerase: promoter structure and polymerase binding. *Biochemistry* 18:528-537.
339. Oen, H., C.-W. Wu, R. Haas, and P. E. Cole. 1979. T7 DNA-directed, rapid-turnover, single-strand addition reactions catalyzed by *E. coli* RNA polymerase. *Biochemistry* 18:4148-4155.
340. Oey, J. L., W. Strätling, and R. Knippers. 1971. A DNA polymerase induced by bacteriophage T7. *Eur. J. Biochem.* 23:497-504.
341. Ogawa, H., H. Araki, and Y. Tsujimoto. 1979. Recombination intermediates formed in the extract from T7-infected cells. *Cold Spring Harbor Symp. Quant. Biol.* 43:1033-1041.
342. Ogawa, T., and T. Okazaki. 1979. RNA-linked nascent DNA pieces in phage T7-infected *E. coli*. III. Detection of intact primer RNA. *Nucleic Acids Res.* 6:1621-1633.
343. Okazaki, R., A. Sugino, S. Hirose, T. Okazaki, Y. Imae, R. Kainuma-Kuroda, T. Ogawa, M. Arisawa, and Y. Kurosawa. 1973. The discontinuous replication of DNA, p. 83-106. In R. D. Wells and R. B. Inman (ed.), *DNA synthesis in vitro*. Medical and Technical Publishing Co., Ltd., Lancaster, England.
344. Ontell, M. P., and D. Nakada. 1980. Rescue of abortive T7 gene 2 mutant phage infection by rifampin. *J. Virol.* 34:438-445.
345. Pahl, C. A., and E. T. Young. 1976. Detection of polycistronic and overlapping bacteriophage T7 late transcripts by in vitro translation. *Proc. Natl. Acad. Sci. U.S.A.* 73:312-316.
346. Pahl, C. A., and E. T. Young. 1978. The size and messenger RNA activity of bacteriophage T7 late transcripts synthesized in vivo. *J. Mol. Biol.* 122:69-101.
347. Pacumbaba, R. R., and M. S. Center. 1973. Association of replicating bacteriophage T7 DNA with bacterial membranes. *J. Virol.* 12:855-862.
348. Pacumbaba, R. R., and M. S. Center. 1974. Studies on an endonuclease activity associated with the bacteriophage T7 DNA-membrane complex. *J. Virol.* 14:1380-1387.
349. Pacumbaba, R. R., and M. S. Center. 1975. Partial purification and properties of a bacteriophage T7 inhibitor of the host exonuclease V activity. *J. Virol.* 16:1200-1207.
350. Paetkau, V., L. Langman, R. Bradley, D. Scraba, and R. C. Miller. 1977. Folded, concatenated genomes as replication intermediates of bacteriophage T7 DNA. *J. Virol.* 22:130-141.
351. Paetkau, V., L. Langman, and R. C., Miller. 1975. The origin of nascent single-stranded fragments in replicating T7 DNA. *J. Mol. Biol.* 98:719-737.
352. Pai, S. H., H. J. Rahmsdorf, H. Ponta, M. Hirsch-Kauffmann, P. Herrlich, and M. Schweiger. 1975. Protein kinase of bacteriophage T7. II. Properties, enzyme synthesis in vitro and regulation of enzyme synthesis and activity in vivo. *Eur. J. Biochem.* 55:305-314.
353. Palchaudhuri, S., and W. K. Maas. 1977. Physical mapping of genes on the F plasmid of *Escherichia coli* responsible for inhibition of growth of female-specific bacteriophages. *J. Bacteriol.* 132:740-743.
354. Panayotatos, N., and R. D. Wells. 1979. Identification, cloning, and characterization of three late promoters at 14.6, 14.8 and 15.9% of the T7 DNA. *J. Mol. Biol.* 135:91-109.
355. Panayotatos, N., and R. D. Wells. 1979. Recognition and initiation site for late promoters of phage T7 is a 22-base pair DNA sequence. *Nature (London)* 280:35-39.
356. Panayotatos, N., and R. D. Wells. 1979. Cloning and localization of the in vitro functional origin of replication of bacteriophage T7 DNA. *J. Biol. Chem.* 254:5555-5561.
357. Pao, C. C., and J. F. Speyer. 1973. Order of injection of T7 bacteriophage DNA. *J. Virol.* 11:1024-1026.
358. Pao, C. C., and J. F. Speyer. 1975. Mutants of T7 bacteriophage inhibited by lambda prophage. *Proc. Natl. Acad. Sci. U.S.A.* 72:3642-3646.
359. Parson, K. A., H. R. Warner, D. L. Anderson, and D. P. Snustad. 1973. Analysis of nuclear disruption and binding of intermediates in host DNA breakdown to membranes after infection of *Escherichia coli* with bacteriophages T4 and T7. *J. Virol.* 11:806-809.
360. Peters, G. G., and R. S. Hayward. 1974. Dinucleotide sequences in the regions of T7 DNA coding for termination of early transcription. *Eur. J. Biochem.* 48:199-208.
361. Peters, G. G., and R. S. Hayward. 1974. Transcriptional termination in vitro: the 3'-terminal sequence of coliphage T7 "early" RNA. *Biochem. Biophys. Res. Commun.* 61:809-816.
362. Pfeffer, S. R., S. J. Stahl, and M. J. Cham-

- berlin. 1977. Binding of *E. coli* RNA polymerase to T7 DNA. Displacement of holoenzyme from promoter complexes by heparin. *J. Biol. Chem.* **252**:5403-5407.
363. Pfeifer, M., A. Betcke, C. Pöhlmann, M. Kurth, and D. H. Liescher. 1980. In vitro and in vivo recombination between DNA molecules in optimized  $\text{CaCl}_2$ -dependent transfection systems: cloning of the *ocr* gene, p. 463-471. In S. Zadrzil and J. Sponar (ed.), DNA recombination, interaction, and repair. Pergamon Press, Ltd., Oxford.
  364. Pfennig-Yeh, M., H. Ponta, M. Hirsch-Kauffmann, H. J. Rahmsdorf, P. Herrlich, and M. Schweiger. 1978. Early T7 gene expression. Rates of RNA synthesis and degradation, protein-kinase-dependent termination of transcription, and efficiency of translation. *Mol. Gen. Genet.* **186**:127-140.
  365. Pigiet, V., and R. R. Conley. 1978. Isolation and characterization of phosphothioredoxin from *E. coli*. *J. Biol. Chem.* **253**:1910-1920.
  366. Ponta, H., K.-H. Altendorf, M. Schweiger, M. Hirsch-Kauffmann, M.-L. Pfennig-Yeh, and P. Herrlich. 1976. *E. coli* membranes become permeable to ions following T7-virus-infection. *Mol. Gen. Genet.* **149**:145-150.
  367. Ponta, H., M. Grätzel, M. Pfennig-Yeh, M. Hirsch-Kauffmann, and M. Schweiger. 1977. Membrane alteration induced by T7 virus infection. *FEBS Lett.* **73**:207-209.
  368. Ponta, H., H. J. Rahmsdorf, S. H. Pai, P. Herrlich, and M. Schweiger. 1974. Control of gene expression in bacteriophage T7. Isolation of a new control protein and mechanism of action. *Mol. Gen. Genet.* **134**:29-38.
  369. Ponta, H., H. J. Rahmsdorf, S. H. Pai, M. Hirsch-Kauffmann, P. Herrlich, and M. Schweiger. 1974. Control of gene expression in bacteriophage T7: transcriptional controls. *Mol. Gen. Genet.* **134**:281-297.
  370. Portmann, R., J. M. Sogo, T. Koller, and W. Zillig. 1974. Binding sites of *E. coli* RNA polymerase on T7 DNA as determined by electron microscopy. *FEBS Lett.* **45**:64-67.
  371. Powling, A., and R. Knippers. 1974. Some functions involved in bacteriophage T7 genetic recombination. *Mol. Gen. Genet.* **134**:173-180.
  372. Powling, A., and R. Knippers. 1976. Recombination of bacteriophage T7 in vivo. *Mol. Gen. Genet.* **149**:63-71.
  373. Prehm, P., B. Jann, G. Schmidt, and S. Stirm. 1976. On a bacteriophage T3 and T4 receptor region within the cell wall lipopolysaccharide of *E. coli* B. *J. Mol. Biol.* **101**:277-281.
  374. Prihn, D. 1975. Nucleotide sequence of an RNA polymerase binding site at an early T7 promoter. *Proc. Natl. Acad. Sci. U.S.A.* **72**:784-788.
  375. Prihn, D. 1975. Bacteriophage T7 early promoters: nucleotide sequences of two RNA polymerase binding sites. *J. Mol. Biol.* **99**:419-443.
  376. Priess, H., B. Koller, R. Hess, and H. Delius. 1980. Electron microscopic mapping and sequence analysis of the terminator for the early message of *E. coli* phage T7. *Mol. Gen. Genet.* **178**:27-34.
  377. Rahmsdorf, H. J., P. Herrlich, S. H. Pai, M. Schweiger, and H. G. Wittmann. 1973. Ribosomes after infection with bacteriophage T4 and T7. *Mol. Gen. Genet.* **127**:259-271.
  378. Rahmsdorf, H. J., S. H. Pai, H. Ponta, P. Herrlich, R. Roskoski, M. Schweiger, and F. W. Studier. 1974. Protein kinase induction in *E. coli* by bacteriophage T7. *Proc. Natl. Acad. Sci. U.S.A.* **71**:586-589.
  379. Rao, R. N., and M. G. Pereira. 1975. Behavior of a hybrid  $F'_{ts114lac^+}$ ,  $his^+$  factor (F42-400) in *Klebsiella pneumoniae* M5a1. *J. Bacteriol.* **123**:792-805.
  380. Ratner, D. 1974. The interaction of bacterial and phage proteins with immobilized *E. coli* RNA polymerase. *J. Mol. Biol.* **88**:373-383.
  381. Ravetch, J. V., K. Horiuchi, and N. D. Zinder. 1978. Nucleotide sequence of the recognition site for the restriction-modification enzyme of *E. coli* B. *Proc. Natl. Acad. Sci. U.S.A.* **75**:2266-2270.
  382. Razin, A., D. Goren, and J. Friedman. 1975. Studies on the biological role of DNA methylation: inhibition of methylation and maturation of bacteriophage  $\phi X174$  by nicotinamide. *Nucleic Acids. Res.* **2**:1967-1974.
  383. Reisbig, R. R., A. Young, M. Woody, and R. M. Woody. 1979. Spectroscopic analysis of the interaction of *E. coli* DNA-dependent RNA polymerase with T7 DNA and synthetic polynucleotides. *J. Biol. Chem.* **254**:11208-11272.
  384. Remes, B., and D. Elseviers. 1980. Adenosine 5'-triphosphate leakage does not cause abortive infection of bacteriophage T7 in male *Escherichia coli*. *J. Bacteriol.* **143**:1054-1056.
  385. Reuben, R. C., and M. L. Gifter. 1973. A DNA-binding protein induced by bacteriophage T7. *Proc. Natl. Acad. Sci. U.S.A.* **70**:1846-1850.
  386. Reuben, R. C., and M. L. Gifter. 1974. A DNA-binding protein induced by bacteriophage T7. Purification and properties of the protein. *J. Biol. Chem.* **249**:3843-3850.
  387. Reuter, M., D. H. Krüger, D. Scholz, and H. A. Rosenthal. 1980. Protection of foreign DNA against host-controlled restriction in bacterial cells. II. Protection of pSF2124 plasmid DNA by the bacteriophage T3 and T7 *ocr*<sup>+</sup> gene function. *Z. Allg. Mikrobiol.* **20**:345-354.
  388. Richardson, C. C., L. J. Romano, R. Kolodner, J. E. LeClerc, F. Tamanoi, M. J. Engler, F. B. Dean, and D. S. Richardson. 1979. Replication of bacteriophage T7 DNA by purified proteins. *Cold Spring Harbor Symp. Quant. Biol.* **43**:427-440.
  389. Richter, D., P. Herrlich, and M. Schweiger. 1972. Phage DNA-directed enzyme synthesis in in vitro system from yeast mitochondria. *Nature (London) New Biol.* **238**:74-76.
  390. Ritchie, D. A. 1975. Control of bacteriophage nucleic acid replication. *Symp. Soc. Gen. Microbiol.* **25**:225-252.

391. Ritchie, D. A., C. A. Thomas, L. A. MacHattie, and P. Wensink. 1967. Terminal repetition in bacteriophage T4 development. *J. Mol. Biol.* 23:365-376.
392. Robb, S. M., D. R. Woods, F. T. Robb, and J. K. Struthers. 1977. Rifampicin-resistant mutant supporting bacteriophage growth on stationary phase *Achromobacter* cells. *J. Gen. Virol.* 35:117-123.
393. Roberts, L., R. Sheldon, and P. D. Sadowski. 1978. Genetic recombination of bacteriophage T7. IV. Asymmetry of recombination frequencies caused by polarity of DNA packaging in vitro. *Virology* 89:252-261.
394. Roberts, R. J. 1976. Restriction endonucleases. *Crit. Rev. Biochem.* 4:123-164.
395. Roberts, R. J. 1977. The role of restriction endonucleases in genetic engineering, p. 21-32. In R. F. Beers and E. G. Bassett (ed.), *Recombinant molecules: impact on science and society*. Raven Press, Publishers, New York.
396. Roberts, R. J. 1978. Restriction endonucleases: a new role in vivo? *Nature (London)* 271:502.
397. Robertson, H. D., and F. Barany. 1979. Enzymes and mechanisms in RNA processing. *FEBS (Fed. Eur. Biochem. Soc.) Proc. Meet.* 51:285-295.
398. Robertson, H. D., E. Dickson, and J. J. Dunn. 1977. A nucleotide sequence from a RNaseIII processing site in bacteriophage T7 RNA. *Proc. Natl. Acad. Sci. U.S.A.* 74:822-826.
399. Roeder, G. S., and P. D. Sadowski. 1979. Pathways of recombination of bacteriophage T7 DNA in vitro. *Cold Spring Harbor Symp. Quant. Biol.* 43:1023-1032.
400. Roeder, S., and P. D. Sadowski. 1977. Bacteriophage T7 morphogenesis. Phage-related particles in cells infected with wild-type and mutant T7 phage. *Virology* 76:263-285.
401. Rohrer, H., and W. Zillig. 1977. Studies on the transcriptional complex of *E. coli* RNA polymerase. *Eur. J. Biochem.* 79:401-409.
402. Romano, L. J., and C. C. Richardson. 1979. Requirements for synthesis of RNA primers during lagging strand synthesis by the DNA polymerase and gene 4 protein of bacteriophage T7. *J. Biol. Chem.* 254:10476-10483.
403. Romano, L. J., and C. C. Richardson. 1979. Characterization of the RNA primers and the DNA product synthesized by the DNA polymerase and gene 4 protein of bacteriophage T7. *J. Biol. Chem.* 254:10483-10489.
404. Rosa, M. D. 1979. Four T7 RNA polymerase promoters contain an identical 23bp sequence. *Cell* 16:815-825.
405. Rosenberg, A. H., M. N. Simon, F. W. Studier, and R. J. Roberts. 1979. Survey and mapping of restriction endonuclease cleavage sites in bacteriophage T7 DNA. *J. Mol. Biol.* 135:907-915.
406. Rosenberg, M., and D. Court. 1979. Regulatory sequences involved in the promotion and termination of RNA transcription. *Annu. Rev. Genet.* 13:319-353.
407. Rosenberg, M., and R. A. Kramer. 1977. Nucleotide sequence surrounding a RNaseIII processing site in bacteriophage T7 RNA. *Proc. Natl. Acad. Sci. U.S.A.* 74:984-988.
408. Rosenberg, M., R. A. Kramer, and J. A. Steitz. 1974. T7 early mRNAs are the direct products of RNaseIII cleavage. *J. Mol. Biol.* 89:777-782.
409. Rosenberg, M., R. A. Kramer, and J. A. Steitz. 1974. The specificity of RNaseIII cleavage of bacteriophage T7 early mRNAs. *Brookhaven Symp. Biol.* 26:277-285.
410. Rothman-Denes, L. B., S. Muthukrishnan, R. Haselkorn, and F. W. Studier. 1973. A T7 gene function required for shutoff of host and early T7 transcription, p. 227-239. In C. F. Fox and W. S. Robinson (ed.), *Virus research*. Academic Press, Inc., New York.
411. Sadowski, P. D. 1972. An endodeoxyribonuclease induced after infection with phage T7 bearing amber mutations in genes 3, 5 and 6. *Can. J. Biochem.* 50:1016-1023.
412. Sadowski, P. D. 1974. Suppression of a mutation in gene 3 of bacteriophage T7 (T7 endonuclease I) by mutations in phage and host polynucleotide ligase. *J. Virol.* 13:226-229.
413. Sadowski, P. D. 1977. Genetic recombination of bacteriophage T7 DNA in vitro. II. Further properties of the in vitro recombination-packaging reaction. *Virology* 78:192-202.
414. Sadowski, P. D., and C. Kerr. 1970. Degradation of *Escherichia coli* B deoxyribonucleic acid after infection with deoxyribonucleic acid-defective amber mutants of bacteriophage T7. *J. Virol.* 6:149-155.
415. Sadowski, P. D., and D. Vetter. 1976. Genetic recombination of bacteriophage T7 DNA in vitro. *Proc. Natl. Acad. Sci. U.S.A.* 73:692-696.
416. Sager, R., and R. Kitchin. 1975. Selective silencing of eukaryotic DNA. *Science* 189:426-433.
417. Saigo, K. 1975. Polar DNA ejection in bacteriophage T7. *Virology* 65:120-127.
418. Saito, H., S. Tabor, F. Tamanai, and C. C. Richardson. 1980. Nucleotide sequence of the primary origin of bacteriophage T7 DNA replication: relationship to adjacent genes and regulatory elements. *Proc. Natl. Acad. Sci. U.S.A.* 77:3917-3921.
419. Sakaki, Y. 1974. Inactivation of the ATP-dependent DNase of *Escherichia coli* after infection with double-stranded DNA phages. *J. Virol.* 14:1611-1612.
420. Sakaki, Y., A. Karu, S. Linn, and H. Echols. 1973. Purification and properties of the gamma protein specified by bacteriophage lambda: an inhibitor of the host *recBC* recombination system. *Proc. Natl. Acad. Sci. U.S.A.* 70:2215-2220.
421. Salser, W. A. 1974. DNA sequencing techniques. *Annu. Rev. Biochem.* 43:923-965.
422. Salvo, R. A., P. R. Chakraborty, and U. Maitra. 1973. Studies on T3-induced RNA polymerase. IV. Transcription of denatured DNA preparations by T3 RNA polymerase. *J. Biol. Chem.* 248:6647-6654.

423. Satta, G., E. Debbia, C. Pruzzo, and L. Calegari. 1979. The peculiar behavior of coliphage *P1vir* mutants on restricting hosts. *Microbios* 22:93-102.
424. Satta, G., C. Pruzzo, E. Debbia, and L. Calegari. 1978. Lysogenic conversion in *Klebsiella pneumoniae*: system which requires active immunity regulation for expression of the conversion phenomenon. *J. Virol.* 28:786-794.
425. Sauerbier, W., M. Schweiger, and P. Herrlich. 1971. Control of gene function in bacteriophage T7. III. Preventing the shutoff of early enzyme synthesis. *J. Virol.* 8:613-618.
426. Schäfer, R., R. Krämer, W. Zillig, and H. Cudny. 1973. On the initiation of transcription by DNA-dependent RNA polymerase from *E. coli*. *Eur. J. Biochem.* 40:367-373.
427. Schaller, H., C. Gray, and K. Herrmann. 1975. Nucleotide sequence of an RNA polymerase binding site form the DNA of bacteriophage fd. *Proc. Natl. Acad. Sci. U.S.A.* 72:737-741.
428. Schell, J. 1979. Transfer of genes into plants via the Ti-plasmid of *A. tumefaciens*. FEBS (Fed. Eur. Biochem. Soc.) Proc. Meet. 51:521-535.
429. Schell, J., S. W. Glover, K. A. Stacey, P. M. A. Broda, and N. Symonds. 1963. The restriction of phage T3 by certain strains of *E. coli*. *Genet. Res.* 4:483-484.
430. Scherzinger, E., P. Herrlich, and M. Schweiger. 1972. Transcription of T3 and T7 early genes by T3 and T7 RNA polymerases. *Mol. Gen. Genet.* 118:67-77.
431. Scherzinger, E., P. Herrlich, M. Schweiger, and H. Schuster. 1971. The early region of the DNA of bacteriophage T7. *Eur. J. Biochem.* 25:341-348.
432. Scherzinger, E., and G. Klotz. 1975. Studies on bacteriophage T7 DNA synthesis in vitro. II. Reconstitution of the T7 replication system using purified proteins. *Mol. Gen. Genet.* 141:233-249.
433. Scherzinger, E., E. Lanka, and G. Hillenbrand. 1977. Role of bacteriophage T7 DNA primase in the initiation of DNA strand synthesis. *Nucleic Acids Res.* 4:4151-4163.
434. Scherzinger, E., E. Lanka, G. Morelli, D. Seifert, and A. Yuki. 1977. Bacteriophage T7-induced DNA priming protein. A novel enzyme involved in DNA replication. *Eur. J. Biochem.* 72:543-558.
435. Scherzinger, E., H. F. Lauppe, N. Voll, and M. Wanke. 1980. Recombinant plasmids carrying promoters, genes and the origin of DNA replication of the early region of bacteriophage T7. *Nucleic Acids Res.* 8:1287-1305.
436. Scherzinger, E., and F. Litfin. 1974. Initiation of the replication of single-stranded DNA by concerted action of phage T7 RNA and DNA polymerase. *Eur. J. Biochem.* 46:179-187.
437. Scherzinger, E., and F. Litfin. 1974. In vitro studies on the role of phage T7 gene 4 product in DNA replication. *Mol. Gen. Genet.* 135:73-86.
438. Scherzinger, E., F. Litfin, and E. Jost. 1973. Stimulation of T7 DNA polymerase by a new phage-coded protein. *Mol. Gen. Genet.* 123:247-262.
439. Scherzinger, E., and D. Seiffert. 1975. Studies on bacteriophage T7 DNA synthesis in vitro. I. Resolution of the T7 replication system into its components. *Mol. Gen. Genet.* 141:213-232.
440. Schlegel, R. A., and C. A. Thomas. 1972. Some special structural features of intracellular bacteriophage T7 concatemers. *J. Mol. Biol.* 68:319-345.
441. Schleicher, M., and E. K. F. Bautz. 1972. Stability of phage T7 gene I mRNA in *E. coli* cells infected with wild type phage and with gene I amber mutants. *FEBS Lett.* 28:139-141.
442. Scholz, D., S. Hansen, and H. A. Rosenthal. 1973. Untersuchungen an T3-Phagen. IV. Isolierung temperatursensitiver Mutanten des Phagen T3 und ihre partielle Charakterisierung. *Z. Allg. Mikrobiol.* 13:425-437.
443. Scholz, D., C. Meissner, and H. A. Rosenthal. 1979. Nutzung von Gen5- und Gen6-ts-Mutanten des Phagen T3 für die Anzeige von Hemmstoffen der DNS-Synthese. *Z. Allg. Mikrobiol.* 19:653-661.
444. Schroeder, C. 1979. The biological specificity of DNA synthesis, p. 53-63. In A. Graffi et al. (ed.), DNA repair and cancer research. Fortschritte der Onkologie, vol. 4. Akademie-Verlag GmbH, Berlin.
445. Schweiger, M., and L. M. Gold. 1969. Bacteriophage T4 DNA-dependent in vitro synthesis of lysozyme. *Proc. Natl. Acad. Sci. U.S.A.* 63:1351-1358.
446. Schweiger, M., M. Hirsch-Kauffman, H. Ponta, M. Pfennig-Yeh, and P. Herrlich. 1975. Biochemistry of T7 development. FEBS (Fed. Eur. Biochem. Soc.) Proc. Meet. 39:55-68.
447. Seki, T., and T. Okazaki. 1979. RNA-linked nascent DNA pieces in phage T7-infected *E. coli*. II. Primary structure of the RNA portion. *Nucleic Acids Res.* 7:1603-1620.
448. Seroika, K., and W. Wackernagel. 1977. In vivo effects of *recBC* DNase, exonuclease I, and DNA polymerases of *Escherichia coli* on the infectivity of native and single-stranded DNA of bacteriophage T7. *J. Virol.* 21:906-912.
449. Serwer, P., and M. E. Pichler. 1978. Electrophoresis of bacteriophage T7 and T7 capsids in agarose gels. *J. Virol.* 28:917-928.
450. Serwer, P. 1974. Fast sedimenting bacteriophage T7 DNA from T7-infected *E. coli*. *Virology* 59:70-88.
451. Serwer, P. 1974. Complexes between bacteriophage T7 capsids and T7 DNA. *Virology* 59:89-107.
452. Serwer, P. 1976. Internal proteins of bacteriophage T7. *J. Mol. Biol.* 107:271-291.
453. Serwer, P. 1979. Fibrous projections from the core of a bacteriophage T7 procapsid. *J. Supramol. Struct.* 11:321-326.
454. Shinozaki, K., and T. Okazaki. 1977. RNA-linked nascent DNA pieces in T7 phage-infected *E. coli* cells. I. Role of gene 6 exonucle-



- ase in removal of the linked RNA. *Mol. Gen. Genet.* **154**:263-267.
455. Shinozaki, K., and T. Okazaki. 1978. T7 gene 6 exonuclease has an RNase H activity. *Nucleic Acids Res.* **5**:4245-4261.
  456. Siebenlist, U. 1979. Nucleotide sequence of the three major early promoters of bacteriophage T7. *Nucleic Acids Res.* **6**:1895-1907.
  457. Siebenlist, U. 1979. RNA polymerase unwinds an 11-base pair segment of a phage T7 promoter. *Nature (London)* **279**:651-652.
  458. Siebenlist, U., and W. Gilbert. 1980. Contacts between *E. coli* RNA polymerase and an early promoter of phage T7. *Proc. Natl. Acad. Sci. U.S.A.* **77**:122-126.
  459. Siegel, P. J., and M. Schaechter. 1973. The role of the host cell membrane in the replication and morphogenesis of bacteriophage. *Annu. Rev. Microbiol.* **27**:261-282.
  460. Silberstein, S., M. Inouye, and F. W. Studier. 1975. Studies on the role of bacteriophage T7 lysozyme during phage infection. *J. Mol. Biol.* **96**:1-11.
  461. Simmon, V. F., and S. Lederberg. 1972. Degradation of bacteriophage lambda DNA after restriction by *E. coli* K-12. *J. Bacteriol.* **112**:161-169.
  462. Simon, M. N., and F. W. Studier. 1973. Physical mapping of the early region of bacteriophage T7 DNA. *J. Mol. Biol.* **79**:249-265.
  463. Skare, J., E. G. Niles, and W. C. Summers. 1974. Localization of the leftmost initiation site for T7 late transcription in vivo and in vitro. *Biochemistry* **13**:3912-3916.
  464. Skavronskaya, A. G., G. I. Aleshkin, and V. V. Demkin. 1979. Specificity of modification-restriction of bacteriophages P1 and lambda controlled by the R124 plasmid in *E. coli* K12 strains. *Genetika* **15**:1719-1723. (In Russian.)
  465. Skurray, R. A., H. Nagaishi, and A. J. Clark. 1976. Molecular cloning of DNA from F sex factor of *E. coli* K12. *Proc. Natl. Acad. Sci. U.S.A.* **73**:64-68.
  466. Smith, H. O. 1979. Nucleotide sequence specificity of restriction endonucleases. *Science* **205**:455-462.
  467. Smith, H. O., and K. W. Wilcox. 1970. A restriction enzyme from *Hemophilus influenzae*. I. Purification and general properties. *J. Mol. Biol.* **51**:379-391.
  468. Sommer, R., and H. Schaller. 1979. Nucleotide sequence of the recognition site of the B-specific restriction-modification system in *E. coli*. *Mol. Gen. Genet.* **168**:331-335.
  469. Spoerel, N., and P. Herrlich. 1979. Colivirus-T3-coded S-adenosymethionine hydrolase. *Eur. J. Biochem.* **95**:227-233.
  470. Spoerel, N., P. Herrlich, and T. A. Bickle. 1979. A novel bacteriophage defence mechanism: the anti-restriction protein. *Nature (London)* **278**:30-34.
  471. Stahl, S. J., and M. J. Chamberlin. 1977. An expanded transcriptional map of T7 bacteriophage. Reading of minor T7 promoter sites in vitro by *E. coli* RNA polymerase. *J. Mol. Biol.* **112**:577-601.
  472. Stahl, S. J., and M. J. Chamberlin. 1978. Transcription of T7 DNA containing modified nucleotides by bacteriophage T7-specific RNA polymerase. *J. Biol. Chem.* **253**:4951-4959.
  473. Staudenbauer, W. L., E. Scherzinger, and E. Lanka. 1979. Replication of the colicin E1 plasmid in extracts of *E. coli*: uncoupling of leading strand from lagging strand synthesis. *Mol. Gen. Genet.* **197**:113-120.
  474. Steitz, J. A., and R. A. Bryan. 1977. Two ribosome binding sites from the gene 0.3 mRNA of bacteriophage T7. *J. Mol. Biol.* **114**:527-543.
  475. Stent, G. 1963. Molecular biology of bacterial viruses. W. H. Freeman & Co., San Francisco.
  476. Strätling, W., and R. Knippers. 1973. Function and purification of gene 4 protein of phage T7. *Nature (London)* **245**:195-197.
  477. Strätling, W., E. Krause, and R. Knippers. 1973. Fast-sedimenting DNA in bacteriophage T7-infected cells. *Virology* **51**:109-119.
  478. Strome, S., and E. T. Young. 1978. Translational control of the expression of bacteriophage T7 gene 0.3. *J. Mol. Biol.* **125**:75-93.
  479. Strome, S., and E. T. Young. 1980. Chemical and functional quantitation of gene 0.3 messenger RNA during T7 infection. *J. Mol. Biol.* **136**:417-432.
  480. Strome, S., and E. T. Young. 1980. Translational discrimination against bacteriophage T7 gene 0.3 messenger RNA. *J. Mol. Biol.* **136**:433-450.
  481. Strothkamp, R. E., J. L. Oakley, and J. E. Coleman. 1980. Promoter melting of T7 RNA polymerase as detected by single-stranded endonuclease digestion. *Biochemistry* **19**:1074-1080.
  482. Stüber, D., and H. Bujard. 1979. Electron microscopy of DNA: determination of absolute molecular weights and linear density. *Mol. Gen. Genet.* **154**:299-303.
  483. Studier, F. W. 1969. The genetics and physiology of bacteriophage T7. *Virology* **39**:562-574.
  484. Studier, F. W. 1972. Bacteriophage T7. *Science* **176**:367-376.
  485. Studier, F. W. 1973. Genetic analysis of non-essential bacteriophage T7 genes. *J. Mol. Biol.* **79**:227-236.
  486. Studier, F. W. 1973. Analysis of bacteriophage T7 early RNAs and proteins on slab gels. *J. Mol. Biol.* **79**:237-248.
  487. Studier, F. W. 1975. Gene 0.3 of bacteriophage T7 acts to overcome the DNA restriction system of the host. *J. Mol. Biol.* **94**:283-295.
  488. Studier, F. W. 1975. Genetic mapping of a mutation that causes ribonuclease III deficiency in *Escherichia coli*. *J. Bacteriol.* **124**:307-316.
  489. Studier, F. W. 1975. Gene expression after bacteriophage T7 infection. *FEBS (Fed. Eur. Biochem. Soc.) Proc. Mett.* **39**:45-53.
  490. Studier, F. W. 1979. Relationships among different strains of T7 and among T7-related bacteriophages. *Virology* **95**:70-84.
  491. Studier, F. W., J. J. Dunn, and E. Buzash-Pollert. 1979. Processing of bacteriophage T7

- RNAs by RNaseIII. Miami Winter Symp. 11: 261-269.
492. Studier, F. W., and R. Hausmann. 1969. Integration of two sets of T7 mutants. *Virology* 39: 587-588.
493. Studier, F. W., and N. R. Movva. 1976. SA-Mase gene of bacteriophage T3 is responsible for overcoming host restriction. *J. Virol.* 19: 136-145.
494. Studier, F. W., A. H. Rosenberg, M. N. Simon, and J. J. Dunn. 1979. Genetic and physical mapping in the early region of bacteriophage T7 DNA. *J. Mol. Biol.* 135:917-937.
495. Sugino, A., C. L. Peebles, K. N. Kreuzer, and N. R. Cozzarelli. 1977. Mechanism of action of nalidixic acid: purification of *E. coli* *nalA* gene product and its relationship to DNA gyrase and a novel nicking-closing enzyme. *Proc. Natl. Acad. Sci. U.S.A.* 74:4767-4771.
496. Summers, W. C. 1972. Regulation of RNA metabolism of T7 and related phages. *Annu. Rev. Genet.* 6:191-203.
497. Summers, W. C., I. Brunovskis, and R. W. Hyman. 1973. The process of infection with coliphage T7. VII. Characterization and mapping of the major in vivo transcription products of the early region. *J. Mol. Biol.* 74:291-300.
498. Summers, W. C., and R. B. Siegel. 1970. Transcription of late phage RNA by T7 RNA polymerase. *Nature (London)* 228:1160-1162.
499. Summers, W. C., and W. Szybalski. 1968. Totally asymmetric transcription of coliphage T7 in vivo: correlation with polyG binding sites. *Virology* 34:9-16.
500. Susskind, M. M., and D. Botstein. 1978. Molecular genetics of bacteriophage P22. *Microbiol. Rev.* 42:385-413.
501. Sutter, D., and W. Doerfler. 1980. Methylation of integrated adenovirus type 12 DNA sequences in transformed cells is inversely correlated with viral gene expression. *Proc. Natl. Acad. Sci. U.S.A.* 77:253-256.
502. Takeya, T., and M. Takanami. 1974. Isolation of DNA segments containing promoters from bacteriophage T3 DNA. *Biochemistry* 13: 5388-5394.
503. Tamanai, F., H. Saito, and C. C. Richardson. 1980. Physical mapping of primary and secondary origins of bacteriophage T7 DNA replication. *Proc. Natl. Acad. Sci. U.S.A.* 77:2656-2660.
504. Tanyashin, V. I. 1979. Restriction endonucleases, p. 36-98. In A. A. Bayed (ed.), *Genetic engineering, results in science and technique. Molecular biology series, vol. 12, part 1.* All-Union Institute for Scientific and Technical Information, Moscow. (In Russian.)
505. Thompson, B. J., E. Domingo, and R. C. Warner. 1980. Pseudolysogeny of *Azotobacter* phages. *Virology* 102:267-277.
506. Tjian, R., and R. Losick. 1975. Control of gene expression in sporulating *Bacillus subtilis* by modification of RNA polymerase: evidence for a specific inhibitor of Sigma factor. *FEBS (Fed. Eur. Biochem. Soc.) Proc. Meet.* 38:277-287.
507. Travers, A. 1976. RNA polymerase specificity and the control of growth. *Nature (London)* 263:641-646.
508. Tsujimoto, Y., and H. Ogawa. 1977. Intermediates in genetic recombination of bacteriophage T7 DNA. *J. Mol. Biol.* 109:423-436.
509. Tsujimoto, Y., and H. Ogawa. 1977. EcoRI-sensitive mutation of T7 phage. *Mol. Gen. Genet.* 150:221-223.
510. Tsujimoto, Y., and H. Ogawa. 1978. Intermediates in genetic recombination of bacteriophage T7 DNA. Biological activity and the role of gene 3 and gene 5. *J. Mol. Biol.* 125:255-273.
511. Usdin, E., R. T. Borchardt, and C. R. Creveling (ed.). 1979. *Transmethylation.* Elsevier-North Holland Publishing Co., New York.
512. Venner, H., and H. Reinert. 1973. Possible role of methylated DNA bases for the transcription of the genetic information. *Z. Allg. Mikrobiol.* 13:613-624.
513. Vetter, D., L. Roberts, J. B. Jackowski, and P. D. Sadowski. 1977. Location of the *ss*<sup>-</sup> mutation of bacteriophage T7 in gene 10, the structural gene for the major capsid protein. *J. Virol.* 22:694-701.
514. Vlachopoulou, P. J., and P. D. Sadowski. 1977. Genetic recombination of bacteriophage T7 DNA in vitro. III. A physical assay for recombinant DNA. *Virology* 78:203-215.
515. Vovis, G. F., and S. Lacks. 1977. Complementary action of restriction enzymes endoRDpnI and endoRDpnII on bacteriophage f1 DNA. *J. Mol. Biol.* 115:525-538.
516. Wackernagel, W., and U. Hermanns. 1974. Inhibition of exonuclease V after infection of *E. coli* by bacteriophage T7. *Biochem. Biophys. Res. Commun.* 60:521-527.
517. Wackernagel, W., K. Seroka, B. Dreiseikelmann, and A. Prell. 1977. Infectivity of bacteriophage T7 DNA, p. 185-192. In A. Portolés et al. (ed.), *Modern trends in bacterial transformation and transfection.* Elsevier/North Holland Biomedical Press, Amsterdam.
518. Watanabe, I., T. Sakurai, K. Furunse, and A. Ando. 1979. "Pseudolysogenization" by RNA phage Q $\beta$ . *Microbiol. Immunol.* 23:1077-1083.
519. Watanabe, T., T. Takano, T. Arai, H. Nishida, and S. Sato. 1966. Episome-mediated transfer of drug resistance in *Enterobacteriaceae*. X. Restriction and modification of phages by  $\phi$ -R factors. *J. Bacteriol.* 92:477-486.
520. Watson, J. D. 1972. Origin of concatemeric DNA. *Nature (London)* New Biol. 239:197-201.
521. Weidel, W. 1955. Bakteriophagen. *Fortschr. Bot.* 17:856-882.
522. Wickner, S. H. 1978. DNA replication proteins of *E. coli*. *Annu. Rev. Biochem.* 47:1163-1191.
523. Wiggs, J. L., J. W. Bush, and M. J. Chamberlin. 1979. Utilization of promoter and terminator sites on bacteriophage T7 DNA by RNA polymerases from a variety of bacterial orders. *Cell* 18:97-109.
524. Williams, L., and G. G. Meynell. 1971. Female-specific phages and F-minus strains of *E. coli*

- K12. *Mol. Gen. Genet.* 113:222-227.
525. Williams, R. C., and M. J. Chamberlin. 1977. Electron microscope studies of transient complexes formed between *E. coli* RNA polymerase holoenzyme and T7 DNA. *Proc. Natl. Acad. Sci. U.S.A.* 74:3740-3744.
526. Wintermantel, G. 1974. The effect of U.V. light on formation and synthetic capacity of DNA-membrane complexes from T7-infected cells of *E. coli*. *Int. J. Radiat. Biol.* 25:105-111.
527. Witkin, E. M. 1976. Ultraviolet mutagenesis and inducible DNA repair in *Escherichia coli*. *Bacteriol. Rev.* 40:869-907.
528. Wolfson, J., and D. Dressler. 1972. Regions of single-stranded DNA in the growing points of replicating bacteriophage T7 chromosomes. *Proc. Natl. Acad. Sci. U.S.A.* 69:2682-2686.
529. Wolfson, J., and D. Dressler. 1979. Bacteriophage T7 DNA replication. An electron microscopic study of the growing point and the role of the T7 gene 4 protein in the formation of DNA fragments. *J. Biol. Chem.* 254:10490-10495.
530. Wolfson, J., D. Dressler, and M. Magazin. 1972. Bacteriophage T7 DNA replication: a linear replicating intermediate. *Proc. Natl. Acad. Sci. U.S.A.* 69:499-504.
531. Worcel, A., and E. Burgi. 1974. Properties of a membrane-attached form of the folded chromosome of *E. coli*. *J. Mol. Biol.* 82:91-105.
532. Yamada, Y., and D. Nakada. 1976. Early to late switch in bacteriophage development: no translational discrimination between T7 early mRNA and late mRNA. *J. Mol. Biol.* 100:35-46.
533. Yamada, Y., and D. Nakada. 1976. Translation of T7 RNA in vitro without cleavage by RNase III. *J. Virol.* 18:1155-1159.
534. Yamada, Y., J. Silnutzer, and D. Nakada. 1978. Mutant of *E. coli* which blocks T7 bacteriophage assembly: accumulation of short T7 DNA. *J. Mol. Biol.* 121:95-111.
535. Yamada, Y., J. Silnutzer, and D. Nakada. 1979. Accumulation of bacteriophage T7 head-related particles in an *Escherichia coli* mutant. *J. Virol.* 31:209-219.
536. Yamada, Y., P. A. Whitaker, and D. Nakada. 1974. Early to late switch in bacteriophage T7 development; functional decay of T7 early messenger RNA. *J. Mol. Biol.* 89:293-303.
537. Yamada, Y., P. A. Whitaker, and D. Nakada. 1974. Functional instability of T7 early mRNA. *Nature (London)* 248:335-338.
538. Yamada, Y., P. A. Whitaker, and D. Nakada. 1975. Chemical stability of bacteriophage T7 early mRNA. *J. Virol.* 16:1683-1687.
539. Yamagishi, M., H. Fujisawa, H. Yamagishi, and T. Minagawa. 1980. Purification of gene 6 product of bacteriophage T3 and its role in in vitro packaging. *Virology* 100:382-389.
540. Yang, F., A. L. Montoya, D. L. Merlo, M. H. Drummond, M. D. Chilton, E. W. Nester, and M. P. Gordon. 1980. Foreign DNA sequences in crown gall teratomas and their fate during the loss of the tumorous traits. *Mol. Gen. Genet.* 177:707-714.
541. Young, R. J., and G. Smith. 1973. The end groups of T7 mRNA. *Biochem. Biophys. Res. Commun.* 53:952-959.
542. Yuan, R., and R. Reiser. 1978. Steps in the reaction mechanism of the *E. coli* plasmid P15-specific restriction endonuclease. *J. Mol. Biol.* 122:433-445.
543. Yura, T., and A. Ishihama. 1979. Genetics of bacterial RNA polymerases. *Annu. Rev. Genet.* 13:59-97.
544. Zabrie, S. K., and M. F. Shemyakin. 1979. Blockage of phage T7 DNA injection into *E. coli* cells by bacterial RNA polymerase inhibitors. *Dokl. Akad. Nauk SSSR* 246:475-478. (In Russian.)
545. Zechel, K. 1978. Initiation of DNA synthesis by RNA. *Curr. Top. Microbiol. Immunol.* 82:71-112.
546. Zillig, W., H. Fujiki, W. Blum, D. Janekovic, M. Schweiger, H. J. Rahmsdorf, H. Ponta, and M. Hirsch-Kauffmann. 1975. In vivo and in vitro phosphorylation of DNA-dependent RNA polymerase of *E. coli* by bacteriophage T7-induced protein kinase. *Proc. Natl. Acad. Sci. U.S.A.* 72:2506-2510.
547. Zinkovskaya, G. G., G. D. Berdyshev, and B. F. Vanyushin. 1978. Time-specific decrease and change in nature of DNA methylation in cattle during aging. *Biokhimiya* 43:1883-1892. (In Russian.)
548. Zograff, Y. N., and E. M. Khourgess. 1973. Inhibitor of association between sigma-factor and core RNA polymerase in *E. coli* infected with bacteriophage T2 or T7. *Mol. Biol. (USSR)* 7:707-713. (In Russian.)